

Product datasheet for **RG218516**

LAMC2 (NM_018891) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	LAMC2 (NM_018891) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	LAMC2
Synonyms:	B2T; BM600; CSF; EBR2; EBR2A; LAMB2T; LAMNB2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG218516 representing NM_018891 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGCTGCGCTCTGGCTGGGCTGCTGCCTCTGCTTCTCGCTCCTCTGCCCGCAGCCCGGGCCACCTCCA
GGAGGGAAGTCTGTGATTGCAATGGGAAGTCCAGGCAGTGTATCTTTGATCGGAACTTCACAGACAAAC
TGTAATGGATCCGCTGCCTCACTGCAATGACAACACTGATGGCATTCACTGCGAGAAGTGAAGAAT
GGCTTTACCGGCACAGAGAAAGGGACCGCTGTTGCCCTGCAATTGTAACCTCAAAGGTTCTCTTAGTG
CTCGATGTGACAACCTCCGACGGTGCAGCTGTAACCAGGTGTGACAGGAGCCAGATGCGACCGATGCT
GCCAGGCTCCACATGCTCATGGATGCGGGGTGCACCCAAGACCAGAGACTGCTAGACTCCAAGTGTGAC
TGTGACCCAGCTGGCATCGCAGGGCCCTGTGACCGGGCCGCTGTGTCTGCAAGCCAGCTGTCACTGGAG
AACGCTGTGATAGGTGTGATCAGGTTACTATAATCTGGATGGGGGAACCTGAGGGCTGTACCCAGTG
TTTCTGCTATGGGCATTAGCCAGCTGCCCGAGCTCTGCAGAATACAGTGTCCATAAGATCACTCTACC
TTTCATCAAGATGTTGATGGCTGGAAGGCTGTCCAACGAAATGGGTCTCTGCAAAGCTCCAATGGTCAC
AGCGCCATCAAGATGTGTTAGCTCAGCCCAACGACTAGACCTGTCTATTTGTGGCTCCTGCCAATT
TCTTGGGAATCAACAGGTGAGCTATGGTCAAAGCTGTCTTTGACTACCGTGTGGACAGAGGAGGCAGA
CACCCATCTGCCCATGATGTGATTTGGAAGGTGCTGGTCTACGGATCACAGCTCCCTTGATGCCACTTG
GCAAGACACTGCCTTGTGGGCTCACCAAGACTTACACATTCAGGTTAAATGAGCATCCAAGCAATAATTG
GAGCCCCAGCTGAGTTACTTTGAGTATCGAAGGTTACTGCGGAATCTCACAGCCCTCCGCATCCGAGCT
ACATATGGAGAATACAGTACTGGGTACATTGACAATGTGACCCTGATTTAGCCCGCCCTGTCTCTGGAG
CCCCAGCACCCCTGGGTTGAACAGTGTATATGTCCTGTTGGGTACAAGGGGCAATTCTGCCAGGATTGTGC
TTCTGGCTACAAGAGAGATTGAGGAGACTGGGGCCTTTTGGCACCTGTATTCTTGTAACTGTCAAGGG
GGAGGGGCTGTGATCCAGACACAGGAGATTGTTATTCAGGGGATGAGAATCCTGACATTGAGTGTGCTG
ACTGCCAATTGGTTTACAACGATCCGCACGACCCCGCAGCTGCAAGCCATGTCCTGTGATAACGG
GTTGAGCTGCTCAGTGTGCCGAGACGGAGGAGTGGTGTGCAATAACTGCCCTCCCGGGTACCAGT



[View online »](#)

GCCCCGTGTGAGCTCTGTGCTGATGGCTACTTTGGGGACCCCTTTGGTGAACATGGCCCAGTGAGGCCTT
 GTCAGCCCTGTCAATGCAACAACAATGTGGACCCAGTGCCTCTGGGAATTGTGACCCGGCTGACAGGCAG
 GTGTTTGAAGTGTATCCACAACACAGCCGGCATCTACTGCGACCAGTGCAAAGCAGGCTACTTCGGGGAC
 CCATTGGCTCCCAACCCAGCAGACAAGTGTGAGCTTGAACCTGTAAACCCATGGGCTCAGAGCCTGTAG
 GATGTCGAAGTGTGGCACCTGTGTTTGAAGCCAGGATTTGGTGGCCCCAAGTGTGAGCATGGAGCATT
 CAGCTGTCCAGCTTGTCTATAATCAAGTGAAGATTCAGATGGATCAGTTTATGCAGCAGCTTCAGAGAATG
 GAGGCCCTGATTTCAAAGGCTCAGGGTGGTGTGAGTAGTACCTGATACAGAGCTGGAAGGCAGGATGC
 AGCAGGCTGAGCAGGCCCTTCAGGACATTCTGAGAGATGCCAGATTTTCAGAAAGGTGCTAGCAGATCCCT
 TGGTCTCCAGTTGGCCAAGGTGAGGAGCCAAGAGAACAGCTACCAGAGCCGCCTGGATGACCTCAAGATG
 ACTGTGAAAAGAGTTCGGGCTCTGGGAAGTCAGTACCAGAACCGAGTTCGGGATACTCACAGGCTCATCA
 CTCAGATGCAGCTGAGCCTGGCAGAAAGTGAAGCTTCTTGGAAACACTAACATTCCTGCCTCAGACCA
 CTACGTGGGGCCAAATGGCTTTAAAAGTCTGGCTCAGGAGGCCACAAGATTAGCAGAAAGCCACGTTGAG
 TCAGCCAGTAACATGGAGCAACTGACAAGGAAACTGAGGACTATTCAAACAAGCCCTCTACTGGTGC
 GCAAGGCCCTGCATGAAGGAGTCGGAAGCGGAAGCGGTAGCCCGACGGTGTGTGGTCAAGGGCTTGT
 GGAAAAATTTGGAGAAAACCAAGTCCCTGGCCAGCAGTTGACAAGGGAGGCCACTCAAGCGGAAATTTGAA
 GCAGATAGGTCTTATCAGCACAGTCTCCGCCTCTGGATTCAAGTGTCTCGGCTTCAGGGAGTCAGTGATC
 AGTCTTTTCAGGTGGAAGAAGCAAGAGGATCAAAACAAAAGCGGATTAAGTCTCAAGCCTGGTAACCAG
 GCATATGGATGAGTTCAAGCGTACACAGAAGAATCTGGGAAACTGGAAAGAAGAAGCACAGCAGCTCTTA
 CAGAATGGAAAAGTGGGAGAGAGAAAATCAGATCAGCTGCTTCCCGTGCCAATCTTGTAAAAGCAGAG
 CACAAGAAGCACTGAGTATGGGCAATGCCACTTTTTATGAAGTTGAGAGCATCCTTAAAACTCAGAGA
 GTTTGACCTGCAGGTGGACAACAGAAAAGCAGAAGCTGAAGAAGCCATGAAGAGACTCTCTACATCAGC
 CAGAAGGTTTCAGATGCCAGTGAACAAGCCAGCAAGCAGAAAGGCCCTGGGGAGCGCTGCTGCTGATG
 CACAGAGGGCAAAGAATGGGGCCGGGAGGCCCTGGAAATCTCCAGTGAGATTGAACAGGAGATTGGGAG
 TCTGAACCTTGGAAAGCCAATGTGACAGCAGATGGAGCCTTGGCCATGGAAAAGGGACTGGCCCTCTCTGAAG
 AGTGAGATGAGGGAAGTGAAGGAGAGCTGGAAGGAAGGAGCTGGAGTTTGACACGAATATGGATGCAG
 TACAGATGGTGATTACAGAAGCCAGAAGTTGATACCAGAGCCAAGAAGCTGGGGTTACAATCCAAGA
 CACACTCAACACATTAGACGGCCTCTGCATCTGATGGGTATG

ACGCGTACGCGGCCGCTCGAG – GFP Tag – GTTTAA

Protein Sequence:

>RG218516 representing NM_018891
 Red=Cloning site Green=Tags(s)

MPALWLGCLCFSLLLPAARATSRREVDCNKGSRQCFDRELHRQTGNFRCLNCNDNTDGIHCEKCKN
 GFYRHRERDRCLPCNCNSKGSLSARCDNSGRCSCKPGVTGARCDRCLPGFHMLMDAGCTQDQRLDLSKCD
 CDPAGIAGPCDAGRVCVCKPAVTGERCDRCRSGYNNLDGGNPEGCTQCFCYGHSASCRSSAEYSVHKITST
 FHQDVDGWKAVQRNGSPAKLQWSQRHQDVFSQAQRLDPVYFVAPAKFLGNQQVSYGQSLSFYDVRDRGGR
 HPSAHDVILEGAGLRITAPLMPLGKTLPCGLTKTYTFRLEHPSNNWSPQLSYFEYRRLRNLTLRIRA
 TYGEYSTGYIDNVTLISARPVSGAPAPWVEQCICPVGYKQFCQDCASGYKRDSARLGPFGTCIPNCQCG
 GGACDPDPTGDCYSGDENPDIECADCPIGFYNDPHDPRSCPCPCHNGFSCSVMPETEVEVNCNCPGVTG
 ARCELADGDFGDPFGEHGPVPRPCQPCQNNVDPASGNCDRLTGRCLKCIHNTAGIYCDQCKAGYFGD
 PLAPNPADKCRACNCNPMGSEPVGCRSDGTCVCKPGFGPNCEHGFSCPACYNQVKIQMDQFMQQLQRM
 EALISKAQGGDGVVPDTELEGRMQAEQALQDILRDAQISEGASRSLGLQLAKVRSQENSYSRLDDLKM
 TVERVRLGSQYQNRVRDTHRLITQMLSLAESEASLGNINIPASDHVYVGNPFGKSLAQEATRLAESHVE
 SASNMEQLTRETEDYSKQALSLVRKALHEGVSGSGSPDGAVVQGLVEKLEKTKSLAQQLTREATQAEIE
 ADRSYQHSLRLLDSVSRLLQGVSDQSFQVEEAKRIKQKADSLSSLVTRHMDEFKRTQKNLGNWKEEAQQLL
 QNGKSGREKSDQLLSRANLAKSRAQEALSMGNATFYEVESILKNLREFDLQVDNRKAEAEAMKRLSYIS
 QKVSDASDKTQQAERALGSAADAQRAKNAGAGEALEISSEIEQETGSLNLEANVTADGALAMEKGLASLK
 SEMREVEGELERKELEFDTNMDAVQMVITEAQKVDTRAKNAGVTIQDTLNTLDGLLHLMGM

TRTRPLE – GFP Tag – V

Restriction Sites:

Sgfl-MluI

Cloning Scheme:


ACCN: NM_018891

ORF Size: 3333 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_018891.1](#), [NP_061486.1](#)

RefSeq Size: 4297 bp


RefSeq ORF: 3336 bp

Locus ID: 3918

UniProt ID: [Q13753](#)

Cytogenetics:	1q25.3
Domains:	LamB, EGF_Lam
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	ECM-receptor interaction, Focal adhesion, Pathways in cancer, Small cell lung cancer
Gene Summary:	<p>Laminins, a family of extracellular matrix glycoproteins, are the major noncollagenous constituent of basement membranes. They have been implicated in a wide variety of biological processes including cell adhesion, differentiation, migration, signaling, neurite outgrowth and metastasis. Laminins, composed of 3 non identical chains: laminin alpha, beta and gamma (formerly A, B1, and B2, respectively), have a cruciform structure consisting of 3 short arms, each formed by a different chain, and a long arm composed of all 3 chains. Each laminin chain is a multidomain protein encoded by a distinct gene. Several isoforms of each chain have been described. Different alpha, beta and gamma chain isomers combine to give rise to different heterotrimeric laminin isoforms which are designated by Arabic numerals in the order of their discovery, i.e. alpha1beta1gamma1 heterotrimer is laminin 1. The biological functions of the different chains and trimer molecules are largely unknown, but some of the chains have been shown to differ with respect to their tissue distribution, presumably reflecting diverse functions in vivo. This gene encodes the gamma chain isoform laminin, gamma 2. The gamma 2 chain, formerly thought to be a truncated version of beta chain (B2t), is highly homologous to the gamma 1 chain; however, it lacks domain VI, and domains V, IV and III are shorter. It is expressed in several fetal tissues but differently from gamma 1, and is specifically localized to epithelial cells in skin, lung and kidney. The gamma 2 chain together with alpha 3 and beta 3 chains constitute laminin 5 (earlier known as kalinin), which is an integral part of the anchoring filaments that connect epithelial cells to the underlying basement membrane. The epithelium-specific expression of the gamma 2 chain implied its role as an epithelium attachment molecule, and mutations in this gene have been associated with junctional epidermolysis bullosa, a skin disease characterized by blisters due to disruption of the epidermal-dermal junction. Two transcript variants resulting from alternative splicing of the 3' terminal exon, and encoding different isoforms of gamma 2 chain, have been described. The two variants are differentially expressed in embryonic tissues, however, the biological significance of the two forms is not known. Transcript variants utilizing alternative polyA_signal have also been noted in literature. [provided by RefSeq, Aug 2011]</p>

Product images:



Circular map for RG218516