

Product datasheet for **RG219609**

Factor XIII (F13B) (NM_001994) Human Tagged ORF Clone

Product data:

| | |
|---------------------------|---|
| Product Type: | Expression Plasmids |
| Product Name: | Factor XIII (F13B) (NM_001994) Human Tagged ORF Clone |
| Tag: | TurboGFP |
| Symbol: | F13B |
| Synonyms: | FXIIIB |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-AC-GFP (PS100010) |
| E. coli Selection: | Ampicillin (100 ug/mL) |



[View online »](#)

ORF Nucleotide
Sequence:

>RG219609 representing NM_001994
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGAGGTTGAAAAACCTGACTTTTATCATCATATTGATAATCTCAGGAGAACTCTATGCAGAAGAGAAAC
CCTGTGGTTTTTCCTCATGTGAAAAATGGAAGAATTGCCCAATATTACTATACTTTTAAAAAGCTTTTACTT
TCCAATGAGCATAGACAAAAAATTGTCATTTTTCTGCTTGGCTGGTTATACCACTGAAAAGTGAAGACAA
GAAGAGCAAACCACGTGTACAACAGAAGGCTGGTCTCCAGAGCCAAGGTGCTTCAAAAAATGCACTAAGC
CTGACCTGAGTAATGGTTACATCTCTGATGTAAAGTTATTGTATAAAATCAAGAGAACATGCATTATGG
TTGCGCTTCAGGGTACAAAACCACTGGAGGGAAGGATGAAGAAGTGGTCAATGTCTCTCTGATGGATGG
TCTTCTCAACCACTGTAGGAAAGAACATGAAACGTGTTGGCTCCTGAATTATATAATGGAAATTATT
CCACAACACAGAAAACATTCAAAGTGAAGGACAAAGTACAATACGAATGTGCTACTGGCTACTACACAGC
TGGAGGAAAGAAGACAGAGGAGTGAATGTCTCACATACGGATGGTCTCTCACACCAAAATGTACCAAA
TTAAAGTGCTCTCTTTAAGATTAATTGAAAATGGTTATTTTCATCCTGTAAAGCAAACCTATGAAGAAG
GAGATGTCGTTCAAGTTTTCTGTGCATGAAAATTATTATCTAAGTGGATCTGATTTAATTCATGCTATAA
CTTTGGTTGGTACCCAGAATCTCCTGTATGCGAAGGAAGAAGAAACAGATGTCCTCCTCCACCTCTGCC
ATAAACTCCAAAATTCAAACACATTCACAACCTTATCGTCATGGAGAAAATAGTTCATATAGAATGTGAAC
TTAATTTTGAGATCCATGGGTGAGCAGAAAATACGTTGTGAAGATGGAAAATGGACAGAACCTCCAAAATG
CATTGAAGGACAGGAGAAGGTAGCCTGTGAGGAACCACTTATTGAAAATGGTGCAGCAAAATTTACAC
TCTAAGATTTATTACAATGGGGATAAAGTGACATATGCATGTAAAAGCGGCTACCTTCTCCATGGATCGA
ATGAGATAACTTGAATCGTGGAAAATGGACACTTCTCCTGAGTGTGTTGAAAATAATGAGAATTGTAA
GCATCCTCCTGTTGAATGAATGGGCTGTTGCAGACGGGATATTGGCAAGCTATGCAACAGGATCCTCA
GTGGAATATAGATGCAATGAATATTACTTACTGAGGGATCAAAAATATCTCGTTGCGAACAAGGAAAAT
GGTCATCCCACCTGTTTCTTGGAAACCATGACTGTTAATGTGGATTACATGAACAGAAAATAACATAGA
AATGAAGTGGAAATATGAAGGGAAAGTCTTACATGGAGATTTAATAGATTTTGTATGTAACAGGGATAT
GACTTATCTCCATTAACCCATTGTCTGAATTATCTGTGCAGTGAACAGAGGAGAAGTGAATATCCTT
TATGACTAGAAAAGAATCTAAAGGAATGTGCACATCTCCTCCTTATTAAACATGGAGTCATTATTAG
TTCAACAGTAGACACCTATGAAAATGGCTCTTCCAGTAGAATACAGATGTTTTGATCACCATTTCTAGAA
GGATCTAGGGAGCCTATTGTTTAGATGGAATGTGGACTACACCACCTTGTGTTTAGAGCCATGCACAT
TATCTTTTACTGAAATGAAAAGAATAATTTACTTCTGAAATGGGATTTTGACAACAGACCACATTTT
GCATGGTGAATATATTGAGTTTATTTGTAGAGGAGATACTTATCCAGCTGAATTATATTTACTGGATCT
ATACTTAGAATGCAATGTGACAGAGGGCAGTAAAATATCCAAGATGATTCCAAGACAAAGCACTCTGT
CTTATCAAGAACCCTAAGAACA

ACGCGTACGCGGCCGCTCGAG – GFP Tag – GTTTAA

Protein Sequence: >RG219609 representing NM_001994
 Red=Cloning site Green=Tags(s)


MRLKNLTFIIILIIISGELYAEEKPCGFPHVENGRIAQYYTFFKSFYFPMIDKKLSFFCLAGYTTESGRQ
 EEQTTCTTEGWSPEPRCFKCKTKPDLNNGYISDVKLLYKIQENMHYGCASGYKTTGGKDEEVVQCLSDGW
 SSQPTCRKEHETCLAPELYNGNYSTTQKTFKVKDKVQYECATGYTAGGKKEEVECLTYGWSLTPKCTK
 LKCSSRLRIENGYFHPVKQTYEEGDVVQFFCHENYYLSGSDLIQCYNFGWYEPSPVCEGRRNRCPPPLP
 INSKIQTHSTTYRHGEIVHIECELNFEIHGSAEIRCEDGKWTEPPKCI EGQEKVACEPPFIENGAANLH
 SKIYYNGDKVTYACKSGYLLHGSNEITCNRGKWTLPPECVENNENCKHPPVVMNGAVADGILASYATGSS
 VEYRCNEYLLRGSKISRCEQGWSSPPVCLPCTVNVDYMNRRNIEMKWYEGKVLHGLDIDFVCKQGY
 DLSPLTPLSELSVQCNRGEVKYPLCTRKESKGMCTSPPLIKHGVIISSVDTYENGSSVEYRCDFHHFLE
 GSREAYCLDGMWTTPLCLEPCTLSFTEMEKNLLLLKWFDFNRPHELHGEYIEFICRGDTYPAELYITGS
 ILRMQCDRGQLKYPRCIPRQSTLSYQEPLRT

TRTRPLE - GFP Tag - V


Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



Plasmid Map:



ACCN: NM_001994

ORF Size: 1983 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001994.1](#), [NP_001985.1](#)

| | |
|-------------------|--|
| RefSeq Size: | 2208 bp |
| RefSeq ORF: | 1986 bp |
| Locus ID: | 2165 |
| UniProt ID: | P05160 |
| Cytogenetics: | 1q31.3 |
| Protein Families: | Druggable Genome |
| Protein Pathways: | Complement and coagulation cascades |
| Gene Summary: | <p>This gene encodes coagulation factor XIII B subunit. Coagulation factor XIII is the last zymogen to become activated in the blood coagulation cascade. Plasma factor XIII is a heterotetramer composed of 2 A subunits and 2 B subunits. The A subunits have catalytic function, and the B subunits do not have enzymatic activity and may serve as a plasma carrier molecules. Platelet factor XIII is comprised only of 2 A subunits, which are identical to those of plasma origin. Upon activation by the cleavage of the activation peptide by thrombin and in the presence of calcium ion, the plasma factor XIII dissociates its B subunits and yields the same active enzyme, factor XIIIa, as platelet factor XIII. This enzyme acts as a transglutaminase to catalyze the formation of gamma-glutamyl-epsilon-lysine crosslinking between fibrin molecules, thus stabilizing the fibrin clot. Factor XIII deficiency is classified into two categories: type I deficiency, characterized by the lack of both the A and B subunits; and type II deficiency, characterized by the lack of the A subunit alone. These defects can result in a lifelong bleeding tendency, defective wound healing, and habitual abortion. [provided by RefSeq, Jul 2008]</p> |