

Product datasheet for RG219762

FHL2 (NM_001039492) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: FHL2 (NM_001039492) Human Tagged ORF Clone

Tag: TurboGFP

Symbol: FHL2

Synonyms: AAG11; DRAL; FHL-2; SLIM-3; SLIM3

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >RG219762 representing NM_001039492
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGACTGAGCGCTTTGACTGCCACCATTGCAACGAATCTCTCTTTTGGCAAGAAGTACATCCTGCGGGAGG
AGAGCCCCTACTGCGTGGTGTGCTTTGAGACCCTGTTCGCCAACACCTGCGAGGAGTGTGGGAAGCCCAT
CGGCTGTGACTGCAAGGACTTGTCTTACAAGGACCGGCACTGCATGAAGCCTGTTTCCACTGCTCGCAG
TGCAGAAACTCACTGGTGGACAAGCCCTTTGCTGCCAAGGAGGACCAGCTGCTCTGTACAGACTGCTATT
CCAACGAGTACTCATCCAAGTGCCAGGAATGCAAGAAGACCATCATGCCAGGTACCCGCAAGATGGAGTA
CAAGGGCAGCAGCTGGCATGAGACCTGCTTCATCTGCCACCGCTGCCAGCAGCCCAATTGGAACCAAGAGT
TTCATCCCCAAAGACAATCAGAATTTCTGTGTGCCCTGCTATGAGAAACAACATGCCATGCAGTGCGTTC
AGTGCAACAAAAGCCCATCACCACGGGAGGGGTCACTTACCGGGAGCAGCCCTGGCACAAGGAGTGCTTCGT
GTGCACCGCCTGCAGGAAGCAGCTGTTCTGGGCAGCGCTTCACAGCTCGCGATGACTTTGCCTACTGCCTG
AACTGCTTCTGTGACTTGTATGCCAAGAAGTGTGCTGGGTGCACCAACCCCATCAGCGGACTTGGTGGCA
CAAAATACATCTCCTTTGAGGAACGGCAGTGGCATAACGACTGCTTTAACTGTAAGAAAGTGCTCCCTCTC
ACTGGTGGGGCCTTGCTCACAGAGAGGGACGACATCCTGTGCCCCGACTGTGGGAAAAGACATC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >RG219762 representing NM_001039492

Red=Cloning site Green=Tags(s)


MTERFDCHHCNESLFGKKYILREESPYCVVCFETLFANTCEECGKPIGCDCKDLSYKDRHWHEACFHCSQ CRNSLVDKPFAAKEDQLLCTDCYSNEYSSKCQECKKTIMPGTRKMEYKGSSWHETCFICHRCQQPIGTKS FIPKDNQNFCVPCYEKQHAMQCVQCKKPITTGGVTYREQPWHKECFVCTACRKQLSGQRFTARDDFAYCL NCFCDLYAKKCAGCTNPISGLGGTKYISFEERQWHNDCFNCKKCSLSLVGRGFLTERDDILCPDCGKDI

Restriction Sites:


Sgfl-Mlul

Cloning Scheme:





Plasmid Map:



ACCN: NM_001039492

ORF Size: 837 bp



FHL2 (NM_001039492) Human Tagged ORF Clone - RG219762

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001039492.3</u>

 RefSeq Size:
 1454 bp

 RefSeq ORF:
 840 bp

 Locus ID:
 2274

 UniProt ID:
 Q14192

Cytogenetics: 2q12.2

Protein Families: Druggable Genome

Gene Summary: This gene encodes a member of the four-and-a-half-LIM-only protein family. Family members

contain two highly conserved, tandemly arranged, zinc finger domains with four highly conserved cysteines binding a zinc atom in each zinc finger. This protein is thought to have a role in the assembly of extracellular membranes. Also, this gene is down-regulated during transformation of normal myoblasts to rhabdomyosarcoma cells and the encoded protein may function as a link between presenilin-2 and an intracellular signaling pathway. Multiple alternatively spliced variants encoding different isoforms have been identified. [provided by

RefSeq, Jan 2016]