

# **Product datasheet for RG219935**

## **UBE2C (NM 181803) Human Tagged ORF Clone**

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: UBE2C (NM\_181803) Human Tagged ORF Clone

Tag: TurboGFP

Symbol: UBE2C

Synonyms: dJ447F3.2; UBCH10

Mammalian Cell Neomycin

Selection:

**Vector:** pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >RG219935 representing NM\_181803

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

GATAGTCCCT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG219935 representing NM\_181803

Red=Cloning site Green=Tags(s)

MASQNRDPAATSVAAARKGAEPSGGAARGPVGKRLQQELMTLMNPTLIVP

TRTRPLE - GFP Tag - V

**Restriction Sites:** Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




### **Cloning Scheme:**





#### Plasmid Map:



**ACCN:** NM\_181803

ORF Size: 150 bp

#### UBE2C (NM\_181803) Human Tagged ORF Clone - RG219935

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 181803.1</u>, <u>NP 861519.1</u>

RefSeq Size: 520 bp
RefSeq ORF: 152 bp
Locus ID: 11065
Cytogenetics: 20q13.12

**Protein Families:** Druggable Genome

**Protein Pathways:** Ubiquitin mediated proteolysis

**Gene Summary:** The modification of proteins with ubiquitin is an important cellular mechanism for targeting

abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, ubiquitin-conjugating enzymes, and ubiquitin-protein ligases. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is required for the destruction of mitotic cyclins and for cell cycle progression, and may be involved in cancer progression. Multiple transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene have been defined on chromosomes 4, 14, 15, 18, and 19. [provided by RefSeq, Aug 2013]