

Product datasheet for **RG220189**

Glucocorticoid Receptor (NR3C1) (NM_000176) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Glucocorticoid Receptor (NR3C1) (NM_000176) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Glucocorticoid Receptor
Synonyms:	GCCR; GCR; GCRST; GR; GRL
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>RG220189 representing NM_000176
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTGCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGCATCGCC**

ATGGACTCCAAGAATCATTAACTCCTGGTAGAGAAGAAAACCCAGCAGTGTGCTTGCTCAGGAGAGGG
 GAGATGTGATGGACTTCTATAAAACCCCTAAGAGGAGGAGCTACTGTGAAGTTTTCTGCGTCTTACCCTC
 ACTGGCTGTCGCTTCTCAATCAGACTCCAAGCAGCGAAGACTTTTTGGTTGATTTTCCAAAAGGCTCAGTA
 AGCAATGCGCAGCAGCCAGATCTGTCCAAGCAGTTTCACTCTCAATGGGACTGTATATGGGAGAGACAG
 AAACAAAAGTGATGGAAATGACCTGGGATTCCCACAGCAGGGCCAAATCAGCCTTCTCGGGGAAAC
 AGACTTAAAGCTTTTGAAGAAAGCATTGCAAACCTCAATAGGTCGACCAGTGTCCAGAGAACCCCAAG
 AGTTCAGCATCCACTGCTGTGTCTGCTGCCCCACAGAGAAGGAGTTTCCAAAACCTCACTCTGATGAT
 CTTCAGAACAGCAACATTTGAAGGGCCAGACTGGCACCAACGGTGGCAATGTGAAATTGTATACCACAGA
 CCAAAGCACCTTTGACATTTTGCAGGATTTGGAGTTTTCTTCTGGGTCCCAGGTAAAGAGACGAATGAG
 AGTCCTTGAGATCAGACCTGTTGATAGATGAAAACCTGTTTGCTTCTCCTCTGGCGGGAGAAGACGATT
 CATTCTTTTGAAGGAACTCGAATGAGGACTGCAAGCCTCTCATTTTACCGGACACTAAACCCAAAAT
 TAAGGATAATGGAGATCTGGTTTTGTCAAGCCCAGTAATGTAACACTGCCCAAGTGAAAACAGAAAA
 GAAGATTTTCATCGAACTCTGCACCCCTGGGGTAATTAAGCAAGAGAAAAGTGGGCACAGTTTACTGTCAGG
 CAAGCTTCTCGGAGCAAATAAATGGTAATAAAATGTCTGCCATTTCTGTTTATGGTGTGAGTACCTC
 TGGAGGACAGATGTACCACTATGACATGAATACAGCATCCCTTCTCAACAGCAGGATCAGAAGCCTATT
 CTTAATGTCATTCCACCAATCCCGTTGGTCCGAAAATGGAATAGGTGCCAAGGATCTGGAGATGACA
 ACTTGACTTCTCTGGGGACTCTGAACCTCCCTGGTCCGAAACAGTTTTTTCTAATGGCTATTCAAGCCCCAG
 CATGAGACCAGATGTAAGCTCTCCTCCATCCAGCTCCTCAACAGCAACAACAGGACCACCTCCCAAACCTC
 TGCTGGTGTGCTCTGATGAAGCTTCAAGATGTCATTATGGAGTCTTAACTTGTGGAAGCTGTAAGTTT
 TCTTCAAAGAGCAGTGGAAAGGACAGCACAATTACCTATGTGCTGGAAGGAATGATTGCATCATCGATAA
 AATTCGAAGAAAAAACTGCCAGCATGCCGCTATCGAAAATGTCTTCAAGGCTGGAATGAACCTGGAAGCT
 CGAAAAACAAAGAAAAAATAAAAGGAATTCAGCAGGCCACTACAGGAGTCTCACAAGAAACCTCTGAAA
 ATCCTGGTAACAAAACAATAGTTCCTGCAACGTTACCACAACCTACCCCTACCCTGGTGTCACTGTTGGA
 GGTATTGAACCTGAAGTGTATATGCAGGATATGATAGCTCTGTTCCAGACTCAACTTGGAGGATCATG
 ACTACGCTCAACATGTTAGGAGGGCGCAAGTATTGCAGCAGTGAATGGGCAAAGGCAATACCAGGTT
 TCAGGAACCTACACCTGGATGACCAATGACCCTACTGCAGTACTCCTGGATGTTTCTTATGGCATTGTC
 TCTGGGTGGAGATCATATAGACAATCAAGTGCAAACCTGCTGTGTTTTGCTCCTGATCTGATTATTAAT
 GAGCAGAGAATGACTCTACCCTGCATGTACGACCAATGTAACACATGCTGTATGTTTCTCTGAGTTAC
 ACAGGCTTCAGGTATCTTATGAAGAGTATCTCTGTATGAAAACCTTACTGCTTCTCTTTCAGTTCTTAA
 GGACGGTCTGAAGAGCCAAGAGCTATTTGATGAAATTAGAATGACCTACATCAAAGAGCTAGGAAAAAGCC
 ATTGTCAGAGGGAAGGAAACTCCAGCCAGAAGTGGCAGCGGTTTTATCAACTGACAAAACCTTTGGATT
 CTATGCATGAAGTGGTTGAAAATCTCCTTAACTATTGCTTCCAAACATTTTGGATAAGACCATGAGTAT
 TGAATTCCTCCGAGATGTTAGCTGAAATCATCACCATCAGATACCAAAATATTCAAACGGAAATATCAA
 AAACCTTCTGTTTCATCAAAG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA


Protein Sequence: >RG220189 representing NM_000176
Red=Cloning site Green=Tags(s)

```
MDSKESLTPGREENPSSVLAQERGDVMDFYKTLRGGATVKVSASSPSLAVASQSDSKQRLLVDFPKGSV
SNAQQPDL SKAVSL SMGLYMGETETKVMGNDLGFPOQQQISLSSGETDLKLL EESIANLNRSTVSPENPK
SSASTAVSAAPTEKEFPKTHSDVSSEQHLKGQGTNGGNVKLYTDDQSTFDILQDLEFSSGSPGKETNE
SPWRSDLLIDENCLL SPLAGEDDSFLL EGNSEDCKPLILPDTKPKIKDNGDLVLSPPSNVTL PQVKTEK
EDFIELCTPGVIKQEKLGTVYQCASFPGANIIGNKMSAISVHGVSTSGGQMYHYDMNTASLSQQQDQKPI
LNVIPPIPVGSENWNRQCQSGDDNLTSLGTLNFPGRTVFSNGYSSPSMRPDVSSPPSSSSTATTGPPPCL
CLVCSDEASGCHYGVLTCGSCVKVFFKRAVEGQHNYLCAGRNDCIIDKIRRKNCPACRYRKLQAGMNLEA
RKTKKKIKGIQQATTGVSQETSENPGNKTIVPATLPQLTPTLVSLLEVIEPEVLYAGYDSSVPDSTWRIM
TTLNMLGGRQVIAAVKWAKAIPGFRNLHLDQMTLLQYSWMFLMAFALGWRYSRQSSANLLCFAPDLIIN
EQRMTLPCMYDQCKHMLYVSELHRLQVSYEEYLCMKTL LLLSSVPKDGLKSQELFDEIRMTYIKELGKA
IVKREGNSSQNWRFYQLTKLLDSMHEVVENLLNYCFQTFLDKTM SIEFPEMLAEIITNQIPKYSNGNIK
KLLFHQK
```

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_000176


ORF Size: 2331 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_000176.2</u> , <u>NP_000167.1</u>
RefSeq Size:	6784 bp
RefSeq ORF:	2334 bp
Locus ID:	2908
UniProt ID:	<u>P04150</u>
Cytogenetics:	5q31.3
Domains:	HOLI, GCR, zf-C4
Protein Families:	Druggable Genome, Nuclear Hormone Receptor, Transcription Factors
Protein Pathways:	Neuroactive ligand-receptor interaction
Gene Summary:	<p>This gene encodes glucocorticoid receptor, which can function both as a transcription factor that binds to glucocorticoid response elements in the promoters of glucocorticoid responsive genes to activate their transcription, and as a regulator of other transcription factors. This receptor is typically found in the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175). [provided by RefSeq, Feb 2011]</p>

Product images:



Circular map for RG220189