

Product datasheet for **RG220377**

Glucocorticoid Receptor (NR3C1) (NM_001020825) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Glucocorticoid Receptor (NR3C1) (NM_001020825) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Glucocorticoid Receptor
Synonyms:	GCCR; GCR; GCRST; GR; GRL
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

ORF Nucleotide
Sequence:

>RG220377 representing NM_001020825
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGCATCGCC**

ATGGACTCCAAGAATCATTAACTCCTGGTAGAGAAGAAAACCCAGCAGTGTGCTTGCTCAGGAGAGGG
GAGATGTGATGGACTTCTATAAAACCCCTAAGAGGAGGAGCTACTGTGAAGTTTTCTGCGTCTTCACCCCTC
ACTGGCTGTCGCTTCTCAATCAGACTCCAAGCAGCGAAGACTTTTGGTTGATTTTCCAAAAGGCTCAGTA
AGCAATGCGCAGCAGCCAGATCTGTCCAAGCAGTTTCACTCTCAATGGGACTGTATATGGGAGAGACAG
AAACAAAAGTGATGGAAATGACCTGGGATTCCCACAGCAGGGCCAAATCAGCCTTCTCGGGGAAAC
AGACTTAAAGCTTTTGAAGAAAGCATTGCAAACCTCAATAGGTCGACCAGTGTCCAGAGAACCCCAAG
AGTTCAGCATCCACTGCTGTGTCTGCTGCCCCACAGAGAAGGAGTTTCCAAAACCTCACTCTGATGAT
CTTCAGAACAGCAACATTTGAAGGGCCAGACTGGCACCAACGGTGGCAATGTGAAATTGTATACCACAGA
CCAAAGCACCTTTGACATTTTGCAAGATTGGAGTTTTCTCTGGGTCCCAGGTAAAGAGACGAATGAG
AGTCCTTGAGATCAGACCTGTTGATAGATGAAAACCTGTTTGCTTCTCCTCTGGCGGGAGAAGACGATT
CATTCTTTTGAAGGAACTCGAATGAGGACTGCAAGCCTCTCATTACCAGACTAAACCCAAAAT
TAAGGATAATGGAGATCTGGTTTTGTCAAGCCCAGTAATGTAACACTGCCCAAGTGAACACAGAAAA
GAAGATTTTCATCGAATCTGCACCCCTGGGGTAATTAAGCAAGAGAACTGGGCACAGTTTACTGTCAGG
CAAGCTTCTCGGAGCAAATAAATGGTAATAAAATGTCTGCCATTTCTGTTTATGGTGTGAGTACCTC
TGGAGGACAGATGTACCACTATGACATGAATACAGCATCCCTTCTCAACAGCAGGATCAGAAGCCTATT
TTAATGTCAATCCCAATCCCGTTGGTCCGAAAATGGAATAGGTGCCAAGGATCTGGAGATGACA
ACTTGACTTCTCTGGGGACTCTGAATCCCTGGTCCGAAACAGTTTTTTCTAATGGCTATTCAAGCCCCAG
CATGAGACCAGATGTAAGCTCTCCTCCATCCAGCTCCTCAACAGCAACAACAGGACCACCTCCCAAACCTC
TGCTGGTGTGCTCTGATGAAGCTTCAGGATGTCATTATGGAGTCTTAACTTGTGGAAGCTGTAAGTTT
TCTTCAAAGAGCAGTGGAAAGACAGCACAATTACCTATGTGCTGGAAGGAATGATTGCATCATCGATAA
AATTCGAAGAAAAAACTGCCAGCATGCCGCTATCGAAAATGTCTTCAGGCTGGAATGAACCTGGAAGCT
CGAAAAACAAAGAAAAAATAAAGGAATTCAGCAGGCCACTACAGGAGTCTCACAAGAAACCTCTGAAA
ATCCTGGTAACAAAACAATAGTTCTGCAACGTTACCACAACCTACCCCTACCCTGGTGTCACTGTTGGA
GGTTATTGAACCTGAAGTGTATATGCAGGATATGATAGCTCTGTTCCAGACTCAACTTGGAGGATCATG
ACTACGCTCAACATGTTAGGAGGGCGCAAGTATTGCAGCAGTGAATGGGCAAAGGCAATACCAGGTT
TCAGGAACCTACACCTGGATGACCAATGACCCTACTGCAGTACTCCTGGATGTTTCTTATGGCATTTC
TCTGGGTGGAGATCATATAGACAATCAAGTGCAAACCTGCTGTGTTTTGCTCCTGATCTGATTATTAAT
GAGCAGAGAATGACTCTACCCTGCATGTACGACCAATGTAACACATGCTGTATGTTTCTCTGAGTTAC
ACAGGCTTCAGGTATCTTATGAAGAGTATCTCTGTATGAAAACCTTACTGCTTCTCTTTCAGTTCTAA
GGACGGTCTGAAGAGCCAAGAGCTATTTGATGAAATTAGAATGACCTACATCAAAGAGCTAGGAAAAGCC
ATTGTCAAGAGGGAAGGAACTCCAGCCAGAATGGCAGCGGTTTTATCAACTGACAAAACCTTTGGATT
CTATGCATGAAAATGTTATGTGGTAAAACAGAAAGCACATCTCACACATTAATC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG220377 representing NM_001020825
Red=Cloning site Green=Tags(s)

MDSKESLTPGREENPSSVLAQERGDVMDFYKTLRGGATVKVSASSPSLAVASQSDSKQRLLVDFPKGSV
 SNAQQPDL SKAVSL SMGLYMGETETKVMGNDLGFPOQQQISLSSGETDLKLL EESIANLNRSTVSPENPK
 SSASTAVSAAPEKEFPKTHSDVSEQQHLKGQGTNGGNVKLYTTDQSTFDILQDLEFSSGSPGKETNE
 SPWRSDLLIDENCLL SPLAGEDDSFLL EGNSEDCKPLILPDTKPKIKDNGDLVLSPPSNVTL PQVKTEK
 EDFIELCTPGVIKQEKLGTVYQCASFPGANIIGNKMSAISVHGVSTSGGQMYHYDMNTASLSQQDQKPI
 FNVIPPIPVGSENWNRQCQSGDDNLTSLGTLNFPGRTVF SNGYSSPSMRPDVSSPPSSSSTATTGPPPKL
 CLVCSDEASGCHYGVLTCGSCVKVFFKRAVEGQHNYLCAGRNDCIIDKIRRKNCPACRYRKLQAGMNEA
 RKTKKKIKGIQQATTGVSQETSENPGNKTIVPATLPQLTPTLVSLLEVIEPEVLYAGYDSSVPDSTWRIM
 TTLNMLGGRQVIAAVKAWAIPGFRNLHLDDQMTLLQYSWMFLMAFALGWRYSYRQSSANLLCFAPDLIIN
 EQRMTLPCMYDQCKHMLYVSELHRLQVSYEEYLCMKTL LLLSSVPKDGLKSQLFDEIRMTYIKELGKA
 IVKREGNSSQNWRFYQLTKLLDSMHENVMWLKPESTSHTLI

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001020825

ORF Size: 2226 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001020825.2](#)

RefSeq Size: 4154 bp

RefSeq ORF: 2229 bp

Locus ID: 2908

UniProt ID: [P04150](#)

Cytogenetics: 5q31.3


Protein Families: Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

Protein Pathways: Neuroactive ligand-receptor interaction

Gene Summary:

This gene encodes glucocorticoid receptor, which can function both as a transcription factor that binds to glucocorticoid response elements in the promoters of glucocorticoid responsive genes to activate their transcription, and as a regulator of other transcription factors. This receptor is typically found in the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175). [provided by RefSeq, Feb 2011]

Product images:



Circular map for RG220377