

## Product datasheet for **RG221205**

### Gemin 1 (SMN2) (NM\_022877) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Gemin 1 (SMN2) (NM_022877) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	SMN2
Synonyms:	BCD541; C-BCD541; GEMIN1; SMNC; TDRD16B
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG221205 representing NM_022877 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGCGATGAGCAGCGCGCGCAGTGGTGGCGCGTCCCGGAGCAGGAGGATTCGGTGTGTTCCGGCGCG  
GCACAGGCCAGAGCGATGATTCTGACATTTGGGATGATACAGCACTGATAAAAGCATATGATAAAGCTGT  
GGCTTCATTTAAGCATGCTCTAAAGAATGGTGACATTTGTGAACTTCGGGTAACCAAAAACCACCT  
AAAAGAAAACCTGCTAAGAAGAATAAAAGCCAAAAGAAGAATACTGCAGTTCCTTACAACAGTGGAAAG  
TTGGGGACAAATGTTCTGCCATTTGGTCAGAAGACGGTTGCATTTACCCAGCTACCATTGCTTCAATTGA  
TTTTAAGAGAGAAACCTGTGTTGTGGTTTACACTGGATATGGAAATAGAGAGGAGCAAAATCTGTCCGAT  
CTACTTTCCCAATCTGTGAAGTAGCTAATAATATAGAACAGAATGCTCAAGAGAATGAAATGAAAGCC  
AAGTTTCAACAGATGAAAGTGAGAACTCCAGGTCTCCTGGAAATAAATCAGATAACATCAAGCCCAATC  
TGCTCCATGGAACCTTTTCTCCCTCCACCACCCCATGCCAGGGCCAAGACTGGGACCAGGAAAGATA  
ATCCCCCACCACCTCCCATATGTCCAGATTCTTTGATGATGCTGATGCTTTGGGAAGTATGTTAATTT  
CATGGTACATGAGTGGCTATCATACTGGCTATTATGAAATGCTGGCA

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

**Protein Sequence:** >RG221205 representing NM\_022877  
Red=Cloning site Green=Tags(s)

MAMSSGGSGGGVPEQEDSVLFRRRTGQSDSDIWDDTALIKAYDKAVASFHALKNGDICETSGKPKTTP  
 KRKPAKKNKSQKKNTAASLQQWKVGDKCSAIWSEDCIYPATIASIDFKRETCVVVYTYGYNREEQNLS  
 LLSPICEVANNIEQNAQENENESQVSTDESENSRSPGNKSDNIKPKSAPWNSFLPPPPMPGPRLGPGKI  
 IPPPPPICPSLDDADALGSMLISWYMSGYHTGYMEMLA

TRTRPLE - GFP Tag - V


**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**

Cloning sites used for ORF Shutting:



**Plasmid Map:**



**ACCN:** NM\_022877

**ORF Size:** 750 bp

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_022877.1</a> , <a href="#">NP_075015.1</a>
<b>RefSeq Size:</b>	1473 bp
<b>RefSeq ORF:</b>	753 bp
<b>Locus ID:</b>	6607
<b>UniProt ID:</b>	<a href="#">Q16637</a>
<b>Cytogenetics:</b>	5q13.2
<b>Protein Families:</b>	Druggable Genome

**Gene Summary:**

This gene is part of a 500 kb inverted duplication on chromosome 5q13. This duplicated region contains at least four genes and repetitive elements which make it prone to rearrangements and deletions. The repetitiveness and complexity of the sequence have also caused difficulty in determining the organization of this genomic region. The telomeric and centromeric copies of this gene are nearly identical and encode the same protein. While mutations in the telomeric copy are associated with spinal muscular atrophy, mutations in this gene, the centromeric copy, do not lead to disease. This gene may be a modifier of disease caused by mutation in the telomeric copy. The critical sequence difference between the two genes is a single nucleotide in exon 7, which is thought to be an exon splice enhancer. Note that the nine exons of both the telomeric and centromeric copies are designated historically as exon 1, 2a, 2b, and 3-8. It is thought that gene conversion events may involve the two genes, leading to varying copy numbers of each gene. The full length protein encoded by this gene localizes to both the cytoplasm and the nucleus. Within the nucleus, the protein localizes to subnuclear bodies called gems which are found near coiled bodies containing high concentrations of small ribonucleoproteins (snRNPs). This protein forms heteromeric complexes with proteins such as SIP1 and GEMIN4, and also interacts with several proteins known to be involved in the biogenesis of snRNPs, such as hnRNP U protein and the small nucleolar RNA binding protein. Four transcript variants encoding distinct isoforms have been described. [provided by RefSeq, Sep 2008]