

## Product datasheet for **RG222125**

### IL37 (NM\_173203) Human Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** IL37 (NM\_173203) Human Tagged ORF Clone  
**Tag:** TurboGFP  
**Symbol:** IL37  
**Synonyms:** FIL1; FIL1(ZETA); FIL1Z; IL-1F7; IL-1H; IL-1H4; IL-1RP1; IL-23; IL-37; IL1F7; IL1H4; IL1RP1  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-AC-GFP (PS100010)  
**E. coli Selection:** Ampicillin (100 ug/mL)  
**ORF Nucleotide Sequence:** >RG222125 representing NM\_173203  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGCC**

ATGTCCTTTGTGGGGGAGAAGCTCAGGAGTGAAAATGGGCTCTGAGGACTGGGAAAAAGATGAACCCAGT  
 GCTGCTTAGAAGAGATCTTCTTTCATTAGCCTCATCCTTGAGCTCAGCCTCTGCGGAGAAAGGAAGTCC  
 GATTCTCCTGGGGTCTCTAAAGGGGAGTTTTGTCTCTACTGTGACAAGGATAAAGGACAAAGTCATCCA  
 TCCCTTCAGCTGAAGAAGGAGAACTGATGAAGCTGGCTGCCAAAAGGAATCAGCACGCCGCCCTTCA  
 TCTTTTATAGGGCTCAGGTGGGCTCCTGGAACATGCTGGAGTCGGCGGCTCACCCGGATGGTTCATCTG  
 CACCTCCTGCAATTGTAATGAGCCTGTTGGGGTGACAGATAAATTTGAGAACAGGAAACACATTGAATTT  
 TCATTTCAACCAGTTTGCAAAGCTGAAATGAGCCCCAGTGAGGTCAGCGAT

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA

**Protein Sequence:** >RG222125 representing NM\_173203  
 Red=Cloning site Green=Tags(s)

MSFVGENSGVKMGSEDWEKDEPQCCLLEEIFFALASSLSSASAEEKGSPILLGVSKGEFCLYCDKDKGQSH  
 SLQLKKEKLMKLAQKESARRPFIFYRAQVGSWNMLESAHPGWFICTSCNENPVGVTDFENRKHIEF  
 SFQPVCKAEMSPSEVSD


**TRTRPLE** - GFP Tag - V

**Restriction Sites:** Sgfl-MluI



[View online »](#)

**Cloning Scheme:**

**Plasmid Map:**


**ACCN:** NM\_173203

**ORF Size:** 471 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u>NM_173203.1, NP_775295.1</u>
<b>RefSeq Size:</b>	604 bp
<b>RefSeq ORF:</b>	474 bp
<b>Locus ID:</b>	27178
<b>UniProt ID:</b>	<u>Q9NZH6</u>
<b>Cytogenetics:</b>	2q14.1
<b>Protein Families:</b>	Druggable Genome, Secreted Protein
<b>Gene Summary:</b>	The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine can bind to, and may be a ligand for interleukin 18 receptor (IL18R1/IL-1Rrp). This cytokine also binds to interleukin 18 binding protein (IL18BP), an inhibitory binding protein of interleukin 18 (IL18), and subsequently forms a complex with IL18 receptor beta subunit, and through which it inhibits the activity of IL18. This gene along with eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. Five alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]