

Product datasheet for RG222463

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cathepsin B (CTSB) (NM_147782) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Cathepsin B (CTSB) (NM 147782) Human Tagged ORF Clone

Tag: TurboGFP

Symbol: CTSB

Synonyms: APPS; CPSB; RECEUP

Mammalian Cell Neo

Selection:

Neomycin

Vector: pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >RG222463 representing NM_147782

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

 ${\tt TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC}$

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA





Protein Sequence: >RG222463 rep

>RG222463 representing NM_147782 Red=Cloning site Green=Tags(s)

MWQLWASLCCLLVLANARSRPSFHPLSDELVNYVNKRNTTWQAGHNFYNVDMSYLKRLCGTFLGGPKPPQ RVMFTEDLKLPASFDAREQWPQCPTIKEIRDQGSCGSCWAFGAVEAISDRICIHTNAHVSVEVSAEDLLT CCGSMCGDGCNGGYPAEAWNFWTRKGLVSGGLYESHVGCRPYSIPPCEHHVNGSRPPCTGEGDTPKCSKI CEPGYSPTYKQDKHYGYNSYSVSNSEKDIMAEIYKNGPVEGAFSVYSDFLLYKSGVYQHVTGEMMGGHAI RILGWGVENGTPYWLVANSWNTDWGDNGFFKILRGQDHCGIESEVVAGIPRTDQYWEKI


TRTRPLE - GFP Tag - V

Restriction Sites:


Sgfl-Mlul

Cloning Scheme:





Plasmid Map:



ACCN: NM_147782

ORF Size: 1017 bp

Cathepsin B (CTSB) (NM_147782) Human Tagged ORF Clone - RG222463

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 147782.4</u>

 RefSeq Size:
 3871 bp

 RefSeq ORF:
 1020 bp

 Locus ID:
 1508

 UniProt ID:
 P07858

Cytogenetics:

Protein Families: Druggable Genome, Protease

8p23.1

Protein Pathways: Antigen processing and presentation, Lysosome

Gene Summary: This gene encodes a member of the C1 family of peptidases. Alternative splicing of this gene

results in multiple transcript variants. At least one of these variants encodes a preproprotein that is proteolytically processed to generate multiple protein products. These products include the cathepsin B light and heavy chains, which can dimerize to form the double chain form of the enzyme. This enzyme is a lysosomal cysteine protease with both endopeptidase and exopeptidase activity that may play a role in protein turnover. It is also known as amyloid

precursor protein secretase and is involved in the proteolytic processing of amyloid

precursor protein (APP). Incomplete proteolytic processing of APP has been suggested to be a causative factor in Alzheimer's disease, the most common cause of dementia. Overexpression of the encoded protein has been associated with esophageal adenocarcinoma and other tumors. Both Cathepsin B and Cathepsin L are involved in the cleavage of the spike protein from the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) upon its entry to the human host cell. Multiple pseudogenes of this gene have been identified. [provided by

RefSeq, Sep 2020]