

Product datasheet for **RG222619**

Apc11 (ANAPC11) (NM_001002248) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Apc11 (ANAPC11) (NM_001002248) Human Tagged ORF Clone
Tag: TurboGFP
Symbol: ANAPC11
Synonyms: APC11; Apc11p; HSPC214
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >RG222619 representing NM_001002248
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGAAGGTGAAGATTAAGTGCTGGAACGGCGTGGCCACTTGGCTCTGGGTGGCCAACGATGAGAAGTGTG
GCATCTGCAGGATGGCATTAAACGGATGCTGCCCTGACTGCAAGGTGCCCGGCGACGACTGCCCGCTGGT
GTGGGGCCAGTGTCCACTGCTCCACATGCATTGCATCCTCAAGTGGCTGCACGCACAGCAGGTGCAG
CAGCACTGCCCATGTGCCCGCCAGGAATGGAAGTCAAGGAG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG222619 representing NM_001002248
Red=Cloning site Green=Tags(s)
MKVKIKCWNGVATWLWVANDENCGICRMAFNGCCPDCKVPGDDCPLVWGQCSHCFHMHCILKWLHAQQVQ
QHCPMCRQEWKFKE

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI



[View online »](#)

OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001002248.3
RefSeq Size:	870 bp
RefSeq ORF:	255 bp
Locus ID:	51529
UniProt ID:	Q9NYG5
Cytogenetics:	17q25.3
Protein Families:	Druggable Genome
Protein Pathways:	Cell cycle, Oocyte meiosis, Progesterone-mediated oocyte maturation, Ubiquitin mediated proteolysis
Gene Summary:	Together with the cullin protein ANAPC2, constitutes the catalytic component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains. May recruit the E2 ubiquitin-conjugating enzymes to the complex.[UniProtKB/Swiss-Prot Function]