

Product datasheet for **RG224323**

STEAP3 (NM_182915) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	STEAP3 (NM_182915) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	STEAP3
Synonyms:	AHMIO2; dudlin-2; dudulin-2; pHyde; STMP3; TSAP6
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:




ACCN:	NM_182915
ORF Size:	1494 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_182915.3
RefSeq Size:	4285 bp
RefSeq ORF:	1497 bp
Locus ID:	55240
UniProt ID:	Q658P3
Cytogenetics:	2q14.2
Protein Families:	Transmembrane
Protein Pathways:	p53 signaling pathway
Gene Summary:	This gene encodes a multipass membrane protein that functions as an iron transporter. The encoded protein can reduce both iron (Fe ³⁺) and copper (Cu ²⁺) cations. This protein may mediate downstream responses to p53, including promoting apoptosis. Deficiency in this gene can cause anemia. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2015]

Product images:



Circular map for RG224323