

Product datasheet for **RG226550**

Retinoic Acid Receptor alpha (RARA) (NM_001145302) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Retinoic Acid Receptor alpha (RARA) (NM_001145302) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	RARA
Synonyms:	NR1B1; RAR
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG226550 representing NM_001145302 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGCCAGCAACAGCAGCTCCTGCCGACACCTGGGGCGGGCACCTCAATGGGTACCCGGTGCCTCCCT
ACGCCTTCTTCTCCCCCTATGCTGGGTGGACTCTCCCCGCCAGGCGCTCTGACCCTCTCCAGCACCA
GCTTCCAGTTAGTGGATATAGCACACCATCCCCAGCCACTGTGAGAAACGACCGAAACAAGAAGAAG
GAGGTGCCAAGCCCGAGTGCTCTGAGAGCTACACGCTGACGCCGAGGTGGGGGAGCTCATTGAGAAGG
TGGCAAAGCGCACCAGAAACCTTCCCTGCCCTCTGCCAGCTGGGCAAATACACTACGAACAACAGCTC
AGAACAACGTGTCTCTCTGGACATTGACCTCTGGGACAAGTTCACTGAACTCTCCACCAAGTGCATCATT
AAGACTGTGGAGTTCGCCAAGCAGCTGCCCGGCTTCAACCCCTCACCATCGCCGACCAGATCACCTCC
TCAAGGCTGCCTGCCTGGACATCCTGATCCTGCGGATCTGCACGCGGTACACGCCGAGCAGGACACCAT
GACCTTCTCGGACGGGCTGACCCTGAACCGGACCCAGATGCACAACGCTGGCTTCGGCCCCCTCACCGAC
CTGGTCTTTGCCTTCGCCAACCAGCTGCTGCCCTGGAGATGGATGATGCGGAGACGGGGCTGCTCAGCG
CCATCTGCCTCATCTGCGGAGACCGCCAGGACCTGGAGCAGCCGACCGGGTGGACATGCTGCAGGAGCC
GCTGCTGGAGGCGCTAAAGGTCTACGTGCGGAAGCGGAGGCCAGCCGCCCCACATGTTCCCAAGATG
CTAATGAAGATTACTGACCTGCGAAGCATCAGCGCCAAGGGGGCTGAGCGGGTATCAGCTGAAGATGG
AGATCCCGGGCTCCATGCCGCTCTCATCCAGGAAATGTTGGAGAACTCAGAGGGCTGGACACTCTGAG
CGGACAGCCGGGGTGGGGGGCGGACGGGGTGGCCTGGCCCCCGCCAGGCAGCTGTAGCCCCAGC
CTCAGCCCCAGCTCCAACAGAAGCAGCCCGCCACCCACTCCCCG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG226550 representing NM_001145302
 Red=Cloning site Green=Tags(s)

MASNSSSCPTPGGGHLNGYPVPPYAFFFPMLGGLSPPGALTTLQHQLPVSQYSTPSPATVRNDRNKKKK
 EVPKPECSESYLTPEVGELIEKVRKAHQETFPALCQLGKYTTNNSSEQRVSLDIDLWDFSELSTKCI
 KTVEFAKQLPGFTTLTIADQITLLKAACLDILILRICTRYTPEQDTMTFSDGLTLNRTQMHNAGFGPLTD
 LVFAFANQLLPLEMDDAETGLLSAICLICGDRQDLEQPDRVMDLQEPLLEALKVYVRKRRPSRPHMFPKM
 LMKITDLRSISAKGAERVITLKMEIPGSMPLIQEMLENSEGLDTLSGQPGGGGRDGGGLAPPGSCSPS
 LSPSSNRSSPATHSP

TRTRPLE - GFP Tag - V


Restriction Sites:

Sgfl-MluI

Cloning Scheme:



Plasmid Map:



ACCN:

NM_001145302

ORF Size:	1095 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001145302.3
RefSeq Size:	3081 bp
RefSeq ORF:	1098 bp
Locus ID:	5914
UniProt ID:	P10276
Cytogenetics:	17q21.2
Protein Families:	Druggable Genome, Nuclear Hormone Receptor, Transcription Factors
Protein Pathways:	Acute myeloid leukemia, Pathways in cancer
Gene Summary:	This gene represents a nuclear retinoic acid receptor. The encoded protein, retinoic acid receptor alpha, regulates transcription in a ligand-dependent manner. This gene has been implicated in regulation of development, differentiation, apoptosis, granulopoiesis, and transcription of clock genes. Translocations between this locus and several other loci have been associated with acute promyelocytic leukemia. Alternatively spliced transcript variants have been found for this locus.[provided by RefSeq, Sep 2010]