

Product datasheet for **RG227545**

IMPA1 (NM_001144878) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: IMPA1 (NM_001144878) Human Tagged ORF Clone
Tag: TurboGFP
Symbol: IMPA1
Synonyms: IMP; IMPA; MRT59
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >RG227545 representing NM_001144878
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGGGCAGCGCCAGGTCCTGTGCTGCCCGCAGTCGCGGTGCTGGGACAGGTGGCAAAAAGGAAGGTGG
 CCTGGCTCCTACGCTGGAAGGCTGTGACCAGGACAGAAACAGCAGGAAACAGTTCAGGTGTTTATGGCTT
 TGGGAAAATGAAAATATTTGTCAAATATTTTCAGAAGATGGCTGATCCTTGGCAGGAATGCATGGATTAT
 GCAGTAACTCTAGCAAGACAAGCTGGAGAGGTAGTTTGTGAAGCTATAAAAAATGAAATGAATGTTATGC
 TGAAAAGTTCTCCAGTTGATTTGGTAACTGCTACGGACCAAAAAGTTGAAAAATGCTTATCTCTCCAT
 AAAGGAAAAGTATCCATCTCACAGTTTCATTGGTGAAGAATCTGTGGCAGCTGGGAAAAAAGTATCTTA
 ACCGACAACCCACATGGATCATTGACCCTATTGATGGAACAACAACTTTGTACATAGATTTCTTTTGT
 TAGCTGTTTCAATTGGCTTTGCTGTAATAAAAAGATAGAATTTGGAGTTGTGTACAGTTGTGTGGAAGG
 CAAGATGTACACTGCCAGAAAAGGAAAGGTGCCTTTTGTAATGGTCAAAAACCAAGTTTCAACAACAA
 GAAGATATTACCAATCTCTCTTGGTACTGAGTTGGGCTCTCCAGAACACCAGAGACTGTGAGAATGG
 TTCTTTTAATATGAAAAGCTTTTTGCATTCTGTTTCCATGGGATCCGGAGTGTGGAACAGCAGCTGT
 TAATATGTGCCTTGTGGCAACTGGCGGAGCAGATGCATATTATGAAATGGGAATTCAGTCTGGGATGTT
 GCAGGAGCTGGCATTATTTACTGAAGCTGGTGGCGTCTAATGGATGTTACAGGTGGACATTTGATT
 TGATGTCACGAAGAGTAATTGCTGCAAATAATAGAATATTAGCAGAAAGGATAGCTAAAGAAATTCAGGT
 TATACCTTTGCAACGAGACGACGAAGAT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG227545 representing NM_001144878
 Red=Cloning site Green=Tags(s)

MGQRPGPVLPAVAVLGQVAKRKVAWLLRWKAVTRTETAGNSSGVYVGFVKMKIFVKYFQKMADPWQECMDY
 AVTLARQAGEVVCFAIKNEMNMLKSSPVDLVTATDQKVEKMLISSIKEKYPSPHSFIEGESVAAGEKSIL
 TDNPTWIIDPIDGTTNFVHRFPFVAVSIGFVAVNKKIEFGVYVSCVEGKMYTARKGKGAFNGQKLOVSQQ
 EDITKSLLVTELSSRTPETVRMVLNMEKLFVIPVHGIRSVGTAAVNMLVATGGADAYEMGIHCWDV
 AGAGIIVTEAGGVLMDVTGGPFDLMSRRVIAANNRILAERIAKEIQVIPLQRDDED

TRTRPLE - GFP Tag - V


Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



Plasmid Map:



ACCN: NM_001144878

ORF Size: 1008 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001144878.1 , NP_001138350.1
RefSeq Size:	3597 bp
RefSeq ORF:	1011 bp
Locus ID:	3612
UniProt ID:	P29218
Cytogenetics:	8q21.13
Protein Families:	Druggable Genome
Protein Pathways:	Inositol phosphate metabolism, Metabolic pathways, Phosphatidylinositol signaling system
Gene Summary:	This gene encodes an enzyme that dephosphorylates myo-inositol monophosphate to generate free myo-inositol, a precursor of phosphatidylinositol, and is therefore an important modulator of intracellular signal transduction via the production of the second messengers myoinositol 1,4,5-trisphosphate and diacylglycerol. This enzyme can also use myo-inositol-1,3-diphosphate, myo-inositol-1,4-diphosphate, scyllo-inositol-phosphate, glucose-1-phosphate, glucose-6-phosphate, fructose-1-phosphate, beta-glycerophosphate, and 2'-AMP as substrates. This enzyme shows magnesium-dependent phosphatase activity and is inhibited by therapeutic concentrations of lithium. Inhibition of inositol monophosphate hydrolysis and subsequent depletion of inositol for phosphatidylinositol synthesis may explain the anti-manic and anti-depressive effects of lithium administered to treat bipolar disorder. Alternative splicing results in multiple transcript variants encoding distinct isoforms. A pseudogene of this gene is also present on chromosome 8q21.13. [provided by RefSeq, Dec 2014]