

## Product datasheet for **RG228898**

### ADA2a (TADA2A) (NM\_001166105) Human Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** ADA2a (TADA2A) (NM\_001166105) Human Tagged ORF Clone  
**Tag:** TurboGFP  
**Symbol:** TADA2A  
**Synonyms:** ADA2; ADA2A; hADA2; KL04P; TADA2L  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-AC-GFP (PS100010)  
**E. coli Selection:** Ampicillin (100 ug/mL)  
**ORF Nucleotide Sequence:** >RG228898 representing NM\_001166105  
**Red=Cloning site Blue=ORF Green=Tags(s)**

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGACCGTTTGGGTTCTTTAGCAATGATCCCTCTGATAAGCCACCTTGCCGAGGCTGCTCCTCTACC  
TCATGGAGCCTTATATCAAGTGTGCTGAATGTGGGCCACCTCTTTTTCTCTGCTTGCAGTGTTCAC  
TCGAGGCTTTGAGTACAAGAAACATCAAAGCGATCATACTTATGAAATAATGACTTCAGATTTTCTGTC  
CTTGATCCAGCTGGACTGCTCAAGAAGAAATGGCCCTTTAGAAGCTGTGATGGACTGTGGCTTTGGAA  
ATTGGCAGGATGTAGCCAATCAAATGTGCACCAAGACCAAGGAGGAGTGTGAGAAGCACTATATGAAGCA  
TTTCATCAATAACCCTCTGTTTGCATCTACCCTGCTGAACCTGAAACAAGCAGAGGAAGCAAAAACCTGCT  
GACACAGCCATTCCATTTCACTTACAGATGACCTCCCCGACCTACCTTTGACTCCTTGCTTTCTCGGG  
ACATGGCCGGGTACATGCCAGCTCGAGCAGATTTTATTGAGGAATTTGACAATTATGCAGATGGGACTT  
GAGAGACATTGATTTTGTGAAGATGACTCGGACATTTTACATGCTCTGAAGATGGCTGTGGTAGATATC  
TATCATTCCAGGTTAAAGGAGAGACAAAGACGAAAAAATTATAAGAGACCATGGATTAATCAACCTTA  
GAAAGTTTCAATTAATGGAACGGCGGTATCCCAAGGAGGTCCAGGACCTGTATGAAACAATGAGGCGATT  
TGCAAGAATTGTGGGCCAGTGGAAATGACAAATTCATTGAAAGCCATGCATTGGAATTTGAACTCCGA  
AGGGAATCAAGAGGCTCCAAGAATACAGGACAGCAGGCATTACCAATTTTGTAGTGCCAGAACCTACG  
ATCACCTCAAGAAGACACGGGAGGAAGAGCGCCTTAAACGCACTATGCTCTCAGAAGTTCTCCAGTATAT  
CCAGGACAGTAGTGCTTGCCAGCAGTGGCTCCGCCGGAAGCTGACATTGATTCCGGCCTGAGTCTTTCC  
ATTCCAATGGCTTCAATTCAGGTAGACGGAGTGCACCACCTTGAACCTCACTGGCCTCCCTGGCACAG  
AGAAGCTGAATGAAAAAGAAAAGGAGCTCTGTCAGATGGTGGTGGTCCCTGGAGCCTATTTAGAATA  
CAAATCTGCTCTATTGAACGAATGTAACAAGCAAGGAGGCTTAAGACTGGCGCAGGCAAGAGCACTCATC  
AAGATAGATGTGAACAAAACCCGAAAACTATGATTTCTCATCAGAGAAGGATACATCACTAAAGGC

**ACGCGTACGCGGCCGCTCGAG** - GFP Tag - GTTTAA



[View online »](#)

**Protein Sequence:** >RG228898 representing NM\_001166105  
 Red=Cloning site Green=Tags(s)

MDRLGSFSNDPSDKPPCRGCSSYLMEPYIKCAECGPPPFLLCLQCFTRGFEYKHKHSDHTYEIMTSDFPV  
 LDPSWTAQEEMALLEAVMDCGFQGNWQDVANQMCTKTKEECEKHYMKHFINNPLFASTLLNLKQAEAAKTA  
 DTAIPFHSTDDPPRPTFDSLLSRDMAGYMPARADFIEEFDNYAEWDLRDIDFVEDDSDILHALKMAVVDI  
 YHSRLKERQRRKKIIRDHGLINLRKFQLMERRYKPEVDLYETMRRFARIVGPVEHDKFIESHALEFELR  
 REIKRLQEYRTAGITNFCSARTYDHLKKTREEERLKRTMLSEVLQYIQDSSACQQWLRQADIDSLSPS  
 IPMASNSGRRSAPPLNLTGLPGTEKLNKEKELCQMVRLVPGAYLEYKSALLNECNKQGGLRLAQRALI  
 KIDVKNKTKIYDFLIREGYITKG

TRTRPLE - GFP Tag - V


**Restriction Sites:**

SgfI-MluI

**Cloning Scheme:**



**Plasmid Map:**



**ACCN:**

NM\_001166105

<b>ORF Size:</b>	1329 bp
<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001166105.3</a>
<b>RefSeq Size:</b>	1678 bp
<b>RefSeq ORF:</b>	1332 bp
<b>Locus ID:</b>	6871
<b>UniProt ID:</b>	<a href="#">Q75478</a>
<b>Cytogenetics:</b>	17q12
<b>Protein Families:</b>	Transcription Factors
<b>Gene Summary:</b>	Many DNA-binding transcriptional activator proteins enhance the initiation rate of RNA polymerase II-mediated gene transcription by interacting functionally with the general transcription machinery bound at the basal promoter. Adaptor proteins are usually required for this activation, possibly to acetylate and destabilize nucleosomes, thereby relieving chromatin constraints at the promoter. The protein encoded by this gene is a transcriptional activator adaptor and has been found to be part of the PCAF histone acetylase complex. Several alternatively spliced transcript variants encoding different isoforms of this gene have been described, but the full-length nature of some of these variants has not been determined. [provided by RefSeq, Oct 2009]