

Product datasheet for RG229608

HDAC8 (NM 001166422) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: HDAC8 (NM 001166422) Human Tagged ORF Clone

Tag: TurboGFP Symbol: HDAC8

Synonyms: CDA07; CDLS5; HD8; HDACL1; KDAC8; MRXS6; RPD3; WTS

Mammalian Cell Neomycin

Selection:

Vector: pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >RG229608 representing NM_001166422
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

TGGCATCATGCAAAGAAAGAGACGTGTGTGTGTATGTGGCACTTTACAAGGCATTC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG229608 representing NM_001166422

Red=Cloning site Green=Tags(s)

MEEPEEPADSGQSLVPVYIYSPEYVSMCDSLAKIPKRASMVHSLIEAYALHKQMRIVKPKVASMEEMATF HTDAYLQHLQKVSQEGDDDHPDSIEYGLGYDCPATEGIFDYAAAIGGATITAAQCLIDGMCKVAINWSGG

WHHAKKETCVYVALYKAF

TRTRPLE - GFP Tag - V

Restriction Sites: Sgfl-Mlul




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Cloning Scheme:





Plasmid Map:



ACCN: NM 001166422

ORF Size: 474 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.



Cytogenetics:

HDAC8 (NM_001166422) Human Tagged ORF Clone - RG229608

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001166422.2</u>

 RefSeq Size:
 1033 bp

 RefSeq ORF:
 477 bp

 Locus ID:
 55869

 UniProt ID:
 Q9BY41

Protein Families: Druggable Genome, Transcription Factors

Xq13.1

Gene Summary: Histones play a critical role in transcriptional regulation, cell cycle progression, and

developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class I of the histone deacetylase family. It catalyzes the deacetylation of lysine residues in the histone N-terminal tails and represses transcription in large multiprotein complexes with transcriptional co-repressors. Multiple transcript variants encoding different isoforms have

been found for this gene. [provided by RefSeq, Oct 2009]