

## **Product datasheet for RG231175**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Caspase 12 (CASP12) (NM\_001191016) Human Tagged ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

Product Name: Caspase 12 (CASP12) (NM\_001191016) Human Tagged ORF Clone

Tag: TurboGFP
Symbol: Caspase 12

Synonyms: CASP-12; CASP12P1

Mammalian Cell

Selection:


Neomycin


**Vector:** pCMV6-AC-GFP (PS100010)


E. coli Selection: Ampicillin (100 ug/mL)

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 







**ACCN:** NM\_001191016

ORF Size: 1023 bp





**OTI Disclaimer:** 

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customercom">customercom</a> or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

**OTI Annotation:** 

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

RefSeq:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

11q22.3

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

NM 001191016.1, NP 001177945.1

 RefSeq Size:
 1688 bp

 RefSeq ORF:
 1026 bp

 Locus ID:
 100506742

 UniProt ID:
 Q6UXS9

**Gene Summary:** 


Cytogenetics:

Caspases are cysteine proteases that cleave C-terminal aspartic acid residues on their substrate molecules. This gene is most highly related to members of the ICE subfamily of caspases that process inflammatory cytokines. In rodents, the homolog of this gene mediates apoptosis in response to endoplasmic reticulum stress. However, in humans this gene contains a polymorphism for the presence or absence of a premature stop codon. The majority of human individuals have the premature stop codon and produce a truncated nonfunctional protein. The read-through codon occurs primarily in individuals of African descent and carriers have endotoxin hypo-responsiveness and an increased susceptibility to severe sepsis. Several alternatively spliced transcript variants have been noted for this gene.

[provided by RefSeq, Feb 2011]



# **Product images:**



Circular map for RG231175