

Product datasheet for RG231178

MDH1 (NM_001199111) Human Tagged ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

sion Plasmids
(NM_001199111) Human Tagged ORF Clone
5FP
EIEE88; HEL-S-32; KAR; MDH-s; MDHA; MGC:1375; MOR2
rcin
-AC-GFP (PS100010)
llin (100 ug/mL)
178 representing NM_001199111 oning site Blue=ORF Green=Tags(s)
AATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC ATCGCC
CGCTGCAGCTATTTTCCAAAGGACGTTACGGTGTTTGATAAGGACGATAAGTCTGAACCAATCA TTGTGACTGGAGCAGCTGGTCAAATTGCATATTCACTGCTGTACAGTATTGGAAATGGATCTGT TAAAGATCAGCCTATAATTCTTGTGCTGTTGGATATCACCCCCATGATGGGTGTCCTGGACGGT ATGGAACTGCAAGACTGTGCCCTTCCCCTCCTGAAAGATGTCATCGCAACAGATAAAGAAGACG TCAAAGACCTGGATGTGGCCATTCTTGTGGGCTCCATGCCAAGAAGGGAAGGCATGGAAGAAGAA ACTGAAAGCAAATGTGAAAATCTTCAAATCCCAGGGTGCAGCCTTAGATAAATACGCCAAGAAG AAGGTTATTGTTGTGGGGTAATCCAGCCAATACCAACTGCCTGACTGCTTCCAAGTCAGCTCCAT CCAAGGAGAACTTCAGTTGCTTGACTCGTTTGGATCACAACCGAGCTAAAGCCAACAGCTCCAT


TAAACTTGGTGTGACTGCTAATGATGTAAAGAATGTCATTATCTGGGGAAACCATTCCTCGACTCAGTAT CCAGATGTCAACCATGCCAAGGTGAAATTGCAAGGAAAGGAAGTTGGTGTTTATGAAGCTCTGAAAGATG ACAGCTGGCTCAAGGGAGAATTTGTCACGACTGTGCAGCAGCGTGGCGCTGCTGTCATCAAGGCTCGAAA

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



ACCN:	NM_001199111
ORF Size:	1056 bp
OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.
	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


ORIGENE MDH1 (NM_001199111) Human Tagged ORF Clone – RG231178	
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 001199111.1, NP 001186040.1</u>
RefSeq Size:	1461 bp
RefSeq ORF:	1059 bp
Locus ID:	4190
UniProt ID:	<u>P40925</u>
Cytogenetics:	2p15
Protein Families:	Druggable Genome
Protein Pathways:	Citrate cycle (TCA cycle), Glyoxylate and dicarboxylate metabolism, Metabolic pathways, Pyruvate metabolism
Gene Summary:	This gene encodes an enzyme that catalyzes the NAD/NADH-dependent, reversible oxidation of malate to oxaloacetate in many metabolic pathways, including the citric acid cycle. Two main isozymes are known to exist in eukaryotic cells: one is found in the mitochondrial matrix and the other in the cytoplasm. This gene encodes the cytosolic isozyme, which plays a key role in the malate-aspartate shuttle that allows malate to pass through the mitochondrial membrane to be transformed into oxaloacetate for further cellular processes. Alternatively spliced transcript variants have been found for this gene. A recent study showed that a C- terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is localized in the peroxisomes. Pseudogenes have been identified on chromosomes X and 6. [provided by RefSeq, Feb 2016]

~ \$1/-

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Product images:



Circular map for RG231178

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US