

Product datasheet for RG231563

Bim (BCL2L11) (NM_207003) Human Tagged ORF Clone

Product data:

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Expression Plasmids |
|------------------------------|---|
| Product Name: | Bim (BCL2L11) (NM_207003) Human Tagged ORF Clone |
| Tag: | TurboGFP |
| Symbol: | BCL2L11 |
| Synonyms: | BAM; BIM; BOD |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-AC-GFP (PS100010) |
| E. coli Selection: | Ampicillin (100 ug/mL) |
| ORF Nucleotide Sequence: | <pre>>RG231563 representing NM_207003 Red=Cloning site Blue=ORF Green=Tags(s)</pre> |
| | TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C |
| | ATGGCAAAGCAACCTTCTGATGTAAGTTCTGAGTGTGACCGAGAAGGTAGACAATTGCAGCCTGCGGAGA GGCCTCCCCAGCTCAGACCTGGGGCCCCTACCTCCCTACAGACAG |
| | ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA |
| Protein Sequence: | <pre>>RG231563 representing NM_207003 Red=Cloning site Green=Tags(s)</pre> |
| | MAKQPSDVSSECDREGRQLQPAERPPQLRPGAPTSLQTEPQASMRQAEPADMRPEIWIAQELRRIGDEFN AYYARRLEK |
| | TRTRPLE - GFP Tag - V |
| Restriction Sites: | Sgfl-Mlul |




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Cloning Scheme:



Plasmid Map:



| ACCN: | NM_207003 |
|-----------------|-----------------------------------|
| ORF Size: | 237 bp |
| OTI Disclaimer: | The molecular sequence of this cl |

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

| GRIGENE Bim (B | CL2L11) (NM_207003) Human Tagged ORF Clone – RG231563 |
|------------------------|---|
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | <u>NM 207003.3</u> |
| RefSeq Size: | 4961 bp |
| RefSeq ORF: | 240 bp |
| Locus ID: | 10018 |
| UniProt ID: | <u>043521</u> |
| Cytogenetics: | 2q13 |
| Protein Families: | Druggable Genome |
| Gene Summary: | The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The protein encoded by this gene contains a Bcl-2 homology domain 3 (BH3). It has been shown to interact with other members of the BCL-2 protein family and to act as an apoptotic activator. The expression of this gene can be induced by nerve growth factor (NGF), as well as by the forkhead transcription factor FKHR-L1, which suggests a role of this gene in neuronal and lymphocyte apoptosis. Transgenic studies of the mouse counterpart suggested that this gene functions as an essential initiator of apoptosis in thymocyte-negative selection. Several alternatively spliced transcript variants of this gene have been identified. [provided by RefSeq, Jun 2013] |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US