

Product datasheet for **RG231669**

ELOC (NM_001204857) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: ELOC (NM_001204857) Human Tagged ORF Clone
Tag: TurboGFP
Symbol: ELOC
Synonyms: SIII; TCEB1
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >RG231669 representing NM_001204857
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGATGGAGAGGAGAAAACCTATGGTGGCTGTGAAGGACCTGATGCCATGTATGTCAAATTGATATCAT
CTGATGGCCATGAATTTATTGTAAAAAGAGAACATGCATTAACATCAGGCACGATAAAAGCCATGTTGAG
TGGCCAGGTCAAGTTGCTGAGAACGAAACCAATGAGGTCAATTTAGAGAGATACCTTCACATGTGCTA
TCGAAAGTATGCATGTATTTACGTACAAGGTTTCGCTACACTAACAGCTCCACCGAGATTCCTGAATTCC
CAATTGCACCTGAAATTGCACTGGAAGTCTGATGGCTGCGAAGTCTTAGATTGT

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG231669 representing NM_001204857
Red=Cloning site Green=Tags(s)
MDGEEKTYGGCEGPDAMYVKLISSDGHEFIVKREHALTSGTIKAMLSGPGQFAENETNEVNFREIPSHVL
SKVCMYFTYKVRYSSTEIPEFPPIAPEIALELLMAANFLDC

TRTRPLE - GFP Tag - V

Restriction Sites: Sgfl-MluI




[View online »](#)

Cloning Scheme:



Plasmid Map:



ACCN: NM_001204857

ORF Size: 336 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001204857.1</u> , <u>NP_001191786.1</u>
RefSeq Size:	1924 bp
RefSeq ORF:	339 bp
Locus ID:	6921
UniProt ID:	<u>Q15369</u>
Cytogenetics:	8q21.11
Protein Families:	Druggable Genome, Transcription Factors
Protein Pathways:	Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysis
Gene Summary:	This gene encodes the protein elongin C, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Multiple alternatively spliced transcript variants encoding two distinct isoforms have been identified. [provided by RefSeq, Mar 2011]