

Product datasheet for **RG232540**

SEPTIN5 (NM_001009939) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	SEPTIN5 (NM_001009939) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	SEPTIN5
Synonyms:	CDCREL; CDCREL-1; CDCREL1; H5; HCDCREL-1; PNUTL1; SEPT5
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG232540 representing NM_001009939 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGACTCGTGGCAGCGCCCCAGGACCGCTGGTGGAGCAGCTGCTGTCGCCGCGGACCCAGGCCAGA
GGCGGCTCAAGGACATTGACAAGCAGTACGTGGGCTTCGCCACTGCCCAACCAGGTGCACCGCAAGTC
GGTGAAGAAAGGCTTTGACTTCACACTCATGGTGGCTGGTGGAGTCAAGGCTGGGGAAGTCCACACTGGTC
CACAGCCTCTTCTGACAGACTTGTACAAGGACCGGAAGCTGCTCAGTGTGAGGAGCGCATCAGCCAGA
CGGTAGAGATTCTAAAACACACGGTGGACATTGAGGAGAAGGGAGTCAAGCTGAAGCTCACCATCGTGGA
CAGCCCGGATTTCGGGGACGCTGTCAACAACACCGAGTGTGGAAGCCATCACCAGACTATGTGGACCAG
CAGTTTGAGCAGTACTTCCGTGATGAGAGCGGCCCAACCGAAAGAATCCAAGACAACCGAGTGCAT
GCTGCCTATACTTCATCTCCCCCTTCGGGCATGGGCTGCGGCCAGTGGATGTGGGTTTCATGAAGGCATT
GCATGAGAAGGTCAACATCGTGCCTCTCATCGCCAAAGCTGACTGTCTTGTCCCCAGTGAGATCCGGAAG
CTGAAGGAGCGGATCCGGGAGGAGATTGACAAGTTTGGGATCCATGTATACCAAGTCCCTGAGTGTGACT
CGGACGAGGATGAGGACTTCAAGCAGCAGGACCGGAACTGAAGGAGAGCGCGCCTTCGCCGTTATAGG
CAGCAACACGGTGGTGGAGGCCAAGGGGACGCGGTCCGGGCGGACTGTACCCCTGGGGATCGTGGAG
GGCGCATTGCGACTTCGTGAAGCTGCGCAACATGCTCATCCGCACGCATATGCACGACCTCAAGGACGTG
ACGTGCGACGTGCACTACGAGAAGTACCGCGCGCACTGCATCCAGCAGATGACCAGCAAAGTACCCAGG
ACAGCCGCATGGAGAGCCCCATCCCGATCCTGCCGCTGCCACCCCGGACGCCGAGAC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG232540 representing NM_001009939
 Red=Cloning site Green=Tags(s)

MDSLAAPQDRLVEQLLSPRTQAQRRLKDIDKQYVGFATLPNQVHRKSVKKGDFDTLMVAGESGLGKSTLV
 HSLFLTDLYKDRKLLSAEERISQTVEILKHTVDIEEKGVKLLTIVDTPGFGDAVNNTECWKPIDYVDQ
 QFEQYFRDESGLNRKNIQDNRVHCCLYFISPFHGRLRPVDVGFMKALHEKVNIPLIAKADCLVPSEIRK
 LKERIREEIDKFGIHVYQFPECDSEDEDFKQDRELKESAPFAVIGSNTVVEAKGQQRVGRGLYPWGIVE
 GALRLREAAQHAHPHAYARPQGRDVRRLRELPRALHPADDQQTDPGPHGEPHPDPAAAHPGRRD

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001009939

ORF Size: 1038 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001009939.3
RefSeq Size:	2284 bp
RefSeq ORF:	1041 bp
Locus ID:	5413
UniProt ID:	Q99719
Cytogenetics:	22q11.21
Protein Families:	Druggable Genome
Protein Pathways:	Parkinson's disease
Gene Summary:	<p>This gene is a member of the septin gene family of nucleotide binding proteins, originally described in yeast as cell division cycle regulatory proteins. Septins are highly conserved in yeast, Drosophila, and mouse and appear to regulate cytoskeletal organization. Disruption of septin function disturbs cytokinesis and results in large multinucleate or polyploid cells. This gene is mapped to 22q11, the region frequently deleted in DiGeorge and velocardiofacial syndromes. A translocation involving the MLL gene and this gene has also been reported in patients with acute myeloid leukemia. Alternative splicing results in multiple transcript variants. The presence of a non-consensus polyA signal (ACAAT) in this gene also results in read-through transcription into the downstream neighboring gene (GP1BB; platelet glycoprotein Ib), whereby larger, non-coding transcripts are produced. [provided by RefSeq, Dec 2010]</p>

Product images:



Circular map for RG232540