

## **Product datasheet for RG233323**

## TAF11 (NM 001270488) Human Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** TAF11 (NM\_001270488) Human Tagged ORF Clone

Tag: TurboGFP

Symbol: TAF11

Synonyms: MGC:15243; PRO2134; TAF2I; TAFII28

Mammalian Cell Neomycin

Selection:

**Vector:** pCMV6-AC-GFP (PS100010)

E. coli Selection: Ampicillin (100 ug/mL)

ORF Nucleotide >RG233323 representing NM\_001270488
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

GTGAGAAGTGGGGAGAAATGCCACCACTACAACCCAAACATA

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG233323 representing NM\_001270488

Red=Cloning site Green=Tags(s)

MDDAHESPSDKGGETGESDETAAVPGDPGATDTDGIPEETDGDADVDLKEAAAEEGELESQDVSDLTTVE REDSSLLNPAAKKLKIDTKEKKEKKQKVDEDEIQKMQILVSSFSEEQLNRYEMYRRSAFPKAAIKRHWMC

VRSGEKCHHYNPNI

TRTRPLE - GFP Tag - V

Restriction Sites: Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Cloning Scheme:**





**ACCN:** NM\_001270488

ORF Size: 462 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001270488.1</u>, <u>NP 001257417.1</u>

 RefSeq Size:
 1490 bp

 RefSeq ORF:
 465 bp

 Locus ID:
 6882

 UniProt ID:
 Q15544



Cytogenetics: 6p21.31

**Protein Families:** Transcription Factors


**Protein Pathways:** Basal transcription factors

**Gene Summary:** Initiation of transcription by RNA polymerase II requires the activities of more than 70

polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID),

which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a small subunit of TFIID that is present in all TFIID complexes and interacts with TBP. This subunit also interacts with another small subunit, TAF13, to form a heterodimer with a structure similar to the histone core structure. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2012]

## **Product images:**



Circular map for RG233323