

## Product datasheet for **RG233533**

### UBE3B (NM\_001270450) Human Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** UBE3B (NM\_001270450) Human Tagged ORF Clone  
**Tag:** TurboGFP  
**Symbol:** UBE3B  
**Synonyms:** BPIDS; KOS  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-AC-GFP (PS100010)  
**E. coli Selection:** Ampicillin (100 ug/mL)  
**ORF Nucleotide Sequence:** >RG233533 representing NM\_001270450  
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGTTACCCCTGTCTCAGACCTCGAGAGCATGGTTCATCGATAGAGCCCGTCAGGCACGAGAAGAAAGGC  
TTGTGCAGAAGGAACGGGAGCGGGCAGCTGTTGTGATCCAGGCCCATGTCCGGAGTTTTCTGTCTCGGAG  
TCGACTGCAGAGAGATATCAGGAGAGAGATTGATGACTTTTTTAAAGCAGATGACCCTGAGTCCACTAAA  
AGAAGTGCACCTTTGTATTTTCAAGATTGCCAGGAACTGCTGTTCTATTTCAGAATCAAAGAGGATAATG  
AGAGATTTGAGAAGTTGTGTCGAGCATCCTGAGCAGCATGGATGCTGAGAATGAGCCTAAGGTGTGGTA  
TGTGTCCCTGGCTTGTCTAAGGACCTCACCTCCTTTGGATTCAACAGATCAAGAACATTTTGTGGTAC  
TGCTGTGATTTTCTCAAGCAGCTCAAGCCTGAAATCCTGCAGGACTCCCGACTCATCACCTGTACCTCA  
CGATGCTTGTACCTTCACAGACTTCAACGTGGAAAATTTCTCGGGGAAAAGGTGAAAGTCTTCGACC  
AGCGATGAACCACATTTGTGCAATATAATGGGACATCTCAACCAGCATGGATTTTATTCTGTGCTGCAG  
TGCTGTGATGGGCTGTTTCTGATTTGGTTTCATATGCTCCTCACAAACCCCTGTGAGGTGGTCCGTTG  
GCAGAAGCTGGTATGACTGGCAGTTGTCTCGC

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

**Protein Sequence:** >RG233533 representing NM\_001270450  
 Red=Cloning site Green=Tags(s)

MFTLSQTSRAWFIDRARQAREERLVQKERERAAVVIQAHVRSFLCRSRLQRDIRREIDDFKADDPSTK  
 RSALCIFKIARKLLFLFRIKEDNERFEKLCRSILSSMDAENEPKVWYVSLACSKDLTLLWIQIKNILWY  
 CCDFLKQLKPEILQDSRLITLTLVTFDTSTWKILRGKGESLRPAMNHICANIMGHLNQHGIFYSLQ  
 CCDGLFPDLVSYAPHNNPVRWSVGRSWYDWQLSR

TRTRPLE - GFP Tag - V

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**ACCN:** NM\_001270450

**ORF Size:** 732 bp


**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001270450.1</a> , <a href="#">NP_001257379.1</a>
<b>RefSeq Size:</b>	1146 bp
<b>RefSeq ORF:</b>	735 bp
<b>Locus ID:</b>	89910
<b>UniProt ID:</b>	<a href="#">Q7Z3V4</a>
<b>Cytogenetics:</b>	12q24.11
<b>Protein Families:</b>	Druggable Genome
<b>Protein Pathways:</b>	Ubiquitin mediated proteolysis
<b>Gene Summary:</b>	<p>The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: E1 ubiquitin-activating enzymes, E2 ubiquitin-conjugating enzymes, and E3 ubiquitin-protein ligases. This gene encodes a member of the E3 ubiquitin-conjugating enzyme family which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme and transfers the ubiquitin to the targeted substrates. A HECT (homology to E6-AP C-terminus) domain in the C-terminus of the longer isoform of this protein is the catalytic site of ubiquitin transfer and forms a complex with E2 conjugases. Shorter isoforms of this protein which lack the C-terminal HECT domain are therefore unlikely to bind E2 enzymes. Alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Jul 2012]</p>

Product images:



Circular map for RG233533