

Product datasheet for **RG233548**

Ikaros (IKZF1) (NM_001220772) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Ikaros (IKZF1) (NM_001220772) Human Tagged ORF Clone
Tag: TurboGFP
Symbol: Ikaros
Synonyms: hlk-1; Hs.54452; IK1; IKAROS; LYF1; PRO0758; ZNFN1A1
Mammalian Cell Selection: Neomycin
Vector: pCMV6-AC-GFP (PS100010)
E. coli Selection: Ampicillin (100 ug/mL)
ORF Nucleotide Sequence: >RG233548 representing NM_001220772
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGATGCTGATGAGGGTCAAGACATGTCCAAGTTTCAGGGGACAAGGGCCTGTCCGACACGCCCTACG
 ACAGCAGCGCCAGCTACGAGAAGGAGAACGAAATGATGAAGTCCCACGTGATGGACCAAGCCATCAACAA
 CGCCATCAACTACCTGGGGCCGAGTCCCTGCGCCGCTGGTGCAGACGCCCCGGGCGTTCCGAGGTG
 GTCCCGGTCATCAGCCCGATGTACCAGCTGCACAAGCCGCTCGCGAGGGCACCCCGCGCTCCAACCACT
 CGGCCACGACAGCGCCGTGGAGAACCTGCTGCTCTCCAAGCCAAGTTGGTGCCCTCGGAGCGCGA
 GCGTCCCGGAGCAACAGCTGCCAAGACTCCACGACACCGAGAGCAACAACGAGGAGCAGCGCAGCGGT
 CTCATCTACCTGACCAACCACATCGCCCCGACGCGCGCAACGGGCTGTGCTCAAGGAGGAGCACC
 CCTACGACCTGCTGCGCGCCGCTCCGAGAAGTCCGAGGACGCGCTCCGCGTGGTCCAGCACCAGCGGGGA
 GCAGATGAAGGTGACAAGTGCGAACACTGCCGGGTGCTCTTCTGGATCACGTCATGTACACCATCCAC
 ATGGGCTGCCACGGTTCCGTGATCCTTTGAGTGCAACATGTGCGGCTACCACAGCCAGGACCGGTACG
 AGTTCTCGTCGCACATAACGCGAGGGGAGCACCCTCCACATGAGC

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG233548 representing NM_001220772
Red=Cloning site Green=Tags(s)

MDADEGQDMSQVSGDKGLSDTPYDSSASYEKENEMMKSHVMDQAINNAINYLGAE SLRPLVQTTPGGSEV
 VPVISP MYQLHKPLAEGTPRSNHS AQDSA VENLLLLSKAKLVPSEREASPSNSCQDSTD TESNNEEQRS G
 LIYLTNHIAPHARNGLSLKEEHRA YDLLRAASENSQDALRVVSTSGEQMKVYKCEHCRVLF LDHVMYTIH
 MGCHGFRDPFECNMCGYHSQDRYEFSSHI TRGEHRFHMS

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001220772

ORF Size: 747 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001220772.1](#), [NP_001207701.1](#)

RefSeq Size: 5407 bp

RefSeq ORF: 750 bp


Locus ID: 10320

Cytogenetics: 7p12.2

Protein Families: Druggable Genome, Transcription Factors

Gene Summary: This gene encodes a transcription factor that belongs to the family of zinc-finger DNA-binding proteins associated with chromatin remodeling. The expression of this protein is restricted to the fetal and adult hemo-lymphopoietic system, and it functions as a regulator of lymphocyte differentiation. Several alternatively spliced transcript variants encoding different isoforms have been described for this gene. Most isoforms share a common C-terminal domain, which contains two zinc finger motifs that are required for hetero- or homo-dimerization, and for interactions with other proteins. The isoforms, however, differ in the number of N-terminal zinc finger motifs that bind DNA and in nuclear localization signal presence, resulting in members with and without DNA-binding properties. Only a few isoforms contain the requisite three or more N-terminal zinc motifs that confer high affinity binding to a specific core DNA sequence element in the promoters of target genes. The non-DNA-binding isoforms are largely found in the cytoplasm, and are thought to function as dominant-negative factors. Overexpression of some dominant-negative isoforms have been associated with B-cell malignancies, such as acute lymphoblastic leukemia (ALL). [provided by RefSeq, May 2014]

Product images:



Circular map for RG233548