

## Product datasheet for **RG233568**

### PCGF5 (NM\_001257101) Human Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** PCGF5 (NM\_001257101) Human Tagged ORF Clone  
**Tag:** TurboGFP  
**Symbol:** PCGF5  
**Synonyms:** RNF159  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-AC-GFP (PS100010)  
**E. coli Selection:** Ampicillin (100 ug/mL)  
**ORF Nucleotide Sequence:** >RG233568 representing NM\_001257101  
**Red=Cloning site Blue=ORF Green=Tags(s)**

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGCTACCCAAAGGAAACACTTGGTCAAAGATTTTAATCCTTACATTACCTGCTATATCTGTAAAGGGT  
ATCTGATCAAGCCAACAACAGTGACGGAATGCCTCCATACATTCTGTAAGACTTGATTGTTTCAGCACTT  
TGAAGATAGCAATGATTGCCAAGGTGTGGCAACCAAGTTCATGAGACAAATCCATTAGAAATGTTGAGG  
TTGGACAATACATTAGAGGAAATATATTTAAGCTGGTCCCTGGACTACGAGAACAAGAAGCTTGAGCGTG  
AATCTGAATTTTGAAGAAAAATAGCCTCAAGAAAATGGACAAGATGATACTTCAAAGCTGACAAACC  
GAAAGTAGATGAAGAAGGTGATGAAAATGAAGATGATAAAGATTATCACAGAAGTGACCCACAAATTGCT  
ATCTGTCTAGATTGTTTACGAAATAATGGCAATCAGGGGACAATGTAGTAAAGGGTTTAAATGAAGAAAT  
TCATTGATGTTCTACACGTGTAAGTGTGGAACTATTAATAAATTTCTAAGTTTAAACTAAAACCTTCC  
AAGTTCTATGAGTTGGATGTGCTGTGCAATGGTAAAATATGGGGAAGGATCATACTATGGAAATTCATC  
TACATGACAAGATGGCGACTAAGAGGCGAAAACCTTCGGTGTCTGAACTGCTCAGCTTCGCAAGTCTGCT  
CTCAGGATGGCCCTTTGTATCAGTCATACCCTATGGTACTGCAGTATCGACCAAGAATTGATTTCCGGT

**ACGCGTACGCGGCCGCTCGAG** - GFP Tag - GTTTAA



[View online »](#)

**Protein Sequence:** >RG233568 representing NM\_001257101  
Red=Cloning site Green=Tags(s)

MATQRKHLVKDFNPYITCYICKGYLIKPTTVTECLHTFCKTCIVQHFEDSNDPCRCGNQVHETNPLEMLR  
 LDNTLEEIIIFKLVPLREQELESEFWKKNKPQENGQDDTSKADKPKVDEEGDENEDDKDYHRSDPQIA  
 ICLDCLRNNGQSGDNVVKGLMKKFIRCSTRVTVTGIKKFLSLKLLPSSYELDLVLCNGEIMGKDHTEFI  
 YMTRWRLRGENFRCLNCSASQVCSQDGPLYQSYPMVLQYRPRIDFG

TRTRPLE - GFP Tag - V


**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**

Cloning sites used for ORF Shutting:



**Plasmid Map:**



**ACCN:** NM\_001257101

**ORF Size:** 768 bp

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001257101.1</a> , <a href="#">NP_001244030.1</a>
<b>RefSeq Size:</b>	7057 bp
<b>RefSeq ORF:</b>	771 bp
<b>Locus ID:</b>	84333
<b>UniProt ID:</b>	<a href="#">Q86SE9</a>
<b>Cytogenetics:</b>	10q23.32
<b>Gene Summary:</b>	Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed:26151332). Within the PRC1-like complex, regulates RNF2 ubiquitin ligase activity (PubMed:26151332). Plays a redundant role with PCGF3 as part of a PRC1-like complex that mediates monoubiquitination of histone H2A 'Lys-119' on the X chromosome and is required for normal silencing of one copy of the X chromosome in XX females (By similarity).[UniProtKB/Swiss-Prot Function]