

Product datasheet for **RG233755**

Chk2 (CHEK2) (NM_001257387) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Chk2 (CHEK2) (NM_001257387) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Chk2
Synonyms:	CDS1; CHK2; hCds1; HuCds1; LFS2; PP1425; RAD53
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG233755 representing NM_001257387 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGTCAAAAACCTTTGGAAGTGGTGCCTGTGGAGAGGTAAGCTGGCTTTCGAGAGGAAAACATGTAAGA
AAGTAGCCATAAAGATCATCAGCAAAAGGAAGTTTGCTATTGGTTCAGCAAGAGAGGCAGACCCAGCTCT
CAATGTTGAAACAGAAATAGAAATTTGAAAAAGCTAAATCATCCTTGATCATCAAGATTA AAAACTTT
TTTGATGCAGAAGATTATTATTTGTTTGAATTGATGGAAGGGGAGAGCTGTTGACAAAGTGGTGG
GGAATAAACGCCTGAAAGAAGCTACCTGCAAGCTCTATTTTACCAGATGCTCTTGCTGTGCAGTACCT
TCATGAAAACGGTATTATACACCGTGACTTAAAGCCAGAGAATGTTTTACTGTCATCTCAAGAAGAGGAC
TGTCTTATAAAGATTACTGATTTTGGGCACTCCAAGATTTTGGGAGAGACCTCTCTCATGAGAACCCTTAT
GTGGAACCCACCTACTTGGCGCCTGAAGTCTTGTCTTCTGTTGGGACTGCTGGGTATAACCGTGTCTGT
GGACTGCTGGAGTTTAGGAGTTATTCTTTTATCTGCCTTAGTGGGTATCCACCTTTCTCTGAGCATAGG
ACTCAAGTGTCACTGAAGGATCAGATCACCAGTGGAAAATACAACCTTCAATCCTGAAGTCTGGGCAGAAG
TCTCAGAGAAAGCTCTGGACCTTGCAAGAAGTTGTTGGTAGTGGATCCAAGGCACGTTTTACGACAGA
AGAAGCCTTAAGACACCCGTGGCTTCAGGATGAAGACATGAAGAGAAAGTTTCAAGATCTTCTGTCTGAG
GAAAATGAATCCACAGCTCTACCCAGGTTCTAGCCCAGCCTTCTACTAGTCAAAGCGGCCCGTGAAG
GGGAAGCCGAGGGTGCCGAGACCACAAAGCGCCAGCTGTGTGCTGCTGTGTTG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

Protein Sequence: >RG233755 representing NM_001257387
 Red=Cloning site Green=Tags(s)

MSKTLGSGACGEVKLAFERKTCKKVAIKIISKRFKAIGSAREADPALNVETEIEILKKNHPICIIKIKNF
 FDAEDYYIVLELMEGGELFDKVVGNKRLKEATCKLYFYQMLLAVQYLHENGIIHRDLKPENVLLSSQEED
 CLIKITDFGHSKILGETSLMRTLCGTPTYLAPVLSVGTAGYNRAVDCWSLGVILFICLSGYPPFSEHR
 TQVSLKDQITSGKYNFIPEVWAEVSEKALDLVKKLLVVDPKARFTTEEALRHPWLQDEDMKRKFQDLLSE
 ENESTALPQVLAQPSTSRKRPREGEAEGAETTKRPAVCAAVL

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001257387

ORF Size: 966 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001257387.2](#)

RefSeq Size: 1976 bp

RefSeq ORF: 969 bp

Locus ID: 11200

UniProt ID: [O96017](#)


Cytogenetics: 22q12.1

Protein Families: Druggable Genome, Protein Kinase, Stem cell - Pluripotency

Protein Pathways: Cell cycle, p53 signaling pathway

Gene Summary: In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Also, mutations in this gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors. This nuclear protein is a member of the CDS1 subfamily of serine/threonine protein kinases. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Apr 2012]

Product images:



Circular map for RG233755