

## **Product datasheet for RG235095**

### Lipin 1 (LPIN1) (NM\_001261428) Human Tagged ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

Product Name: Lipin 1 (LPIN1) (NM\_001261428) Human Tagged ORF Clone

Tag: TurboGFP

Symbol: Lipin 1

Synonyms: PAP1

Mammalian Cell Neomycin

Selection:


**Vector:** pCMV6-AC-GFP (PS100010)


E. coli Selection: Ampicillin (100 ug/mL)

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 

Cloning sites used for ORF Shuttling:





**ACCN:** NM\_001261428

ORF Size: 2925 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

#### Lipin 1 (LPIN1) (NM\_001261428) Human Tagged ORF Clone - RG235095

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001261428.3</u>

 RefSeq Size:
 5620 bp

 RefSeq ORF:
 2928 bp

 Locus ID:
 23175

 UniProt ID:
 Q14693

Cytogenetics: 2p25.1

**Gene Summary:** This gene encodes a magnesium-ion-dependent phosphatidic acid phosphohydrolase


enzyme that catalyzes the penultimate step in triglyceride synthesis including the dephosphorylation of phosphatidic acid to yield diacylglycerol. Expression of this gene is required for adipocyte differentiation and it also functions as a nuclear transcriptional coactivator with some peroxisome proliferator-activated receptors to modulate expression of

other genes involved in lipid metabolism. Mutations in this gene are associated with metabolic syndrome, type 2 diabetes, acute recurrent rhabdomyolysis, and autosomal recessive acute recurrent myoglobinuria (ARARM). This gene is also a candidate for several

human lipodystrophy syndromes. [provided by RefSeq, Mar 2017]



# **Product images:**



Circular map for RG235095