

## Product datasheet for **RG236550**

### SLAMF7 (NM\_001282589) Human Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** SLAMF7 (NM\_001282589) Human Tagged ORF Clone  
**Tag:** TurboGFP  
**Symbol:** SLAMF7  
**Synonyms:** 19A; CD319; CRACC; CS1  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-AC-GFP (PS100010)  
**E. coli Selection:** Ampicillin (100 ug/mL)  
**ORF Nucleotide Sequence:** >RG236550 representing NM\_001282589.  
 Blue=ORF Red=Cloning site Green=Tag(s)

```
GCTCGTTTGTAGTAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCCGCATCGCC
ATGGCTGGTTCCCAACATGCCTCACCCCTCATCTATATCCTTTGGCAGCTCACAGGGTCAGCAGCCTCT
GGACCCGTGAAAGAGCTGGTCCGTTCCGTTGGTGGGGCCGTGACTTCCCCCTGAAGTCCAAAGTAAAG
CAAGTTGACTCTATTGTCTGGACCTTCAACACAACCCCTCTTGTACCATACAGCCAGAAGGGGGCACT
ATCATAGTGACCCAAAATCGTAATAGGGAGAGAGTAGACTTCCAGATGGAGGCTACTCCCTGAAGCTC
AGCAAAGTGAAGAAGAATGACTCAGGGATCTACTATGTGGGGATATACAGCTCATCACTCCAGCAGCCC
TCCACCCAGGAGTACGTGCTGCATGTCTACGAGTACATTGAAGAGAAGAAGAGAGTGGACATTTGTCGG
GAAACTCCTAACATATGCCCCATTCTGGAGAGAACACAGAGTACGACACAATCCCTCACACTAATAGA
ACAATCCTAAAGGAAGATCCAGCAAATACGGTTTACTCCACTGTGGAAATACCGAAAAAGATGGAAAAAT
CCCCACTCACTGCTCACGATGCCAGACACCAAGGCTATTTGCCTATGAGAATGTTATC
ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAAAC
```

**Protein Sequence:** >Peptide sequence encoded by RG236550  
 Blue=ORF Red=Cloning site Green=Tag(s)

```
MAGSPTCLTIYILWQLTGSAAAGPVKELVGSVGGAVTFPLKSKVKQVDSIVWTFNTTPLVTIQPEGGT
IIVTQNRNRERVDFFDGGYSLKLSKLLKNDSGIYYVGIYSSSLQQPSTQEYVLHVYIEIEKKRVDICR
ETPNICPHSGENTEYDTPHNTNRTILKEDPANTVYSTVEIPKMMENPHSLLTMPDTPRLFAYENVI
TRTRPLEMESDESGLPAMEIECRITGTLNGVEFELVGGEGTPEQGRMTNKMSTKGALTFSPYLLSHV
MGYGFYHFGTYPYSGYENPFLHAINNGGYNTRIEKYEDGGVLHVSFSYRYEAGRVIGDFKVMGTGFPED
SVIFTDKIIRSNATVEHLHPMGDNDLDGSFTRTFSLRDGGYSSVVDSHMHFKSAIHP SILQNGGPMFA
FRRVEEDHSNTELGIVEYQHAFKTPDADAGEERV
```

**Restriction Sites:** Sgfl-MluI




**Cloning Scheme:**


- ACCN:** NM\_001282589
- ORF Size:** 612 bp
- OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)
- OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
- RefSeq:** [NM\\_001282589.1](#), [NP\\_001269518.1](#)
- RefSeq Size:** 2515 bp
- RefSeq ORF:** 615 bp
- Locus ID:** 57823
- UniProt ID:** [Q9NQ25](#)
- Cytogenetics:** 1q23.3
- Protein Families:** Druggable Genome, Transmembrane
- MW:** 23.2 kDa

**Gene Summary:**

Self-ligand receptor of the signaling lymphocytic activation molecule (SLAM) family. SLAM receptors triggered by homo- or heterotypic cell-cell interactions are modulating the activation and differentiation of a wide variety of immune cells and thus are involved in the regulation and interconnection of both innate and adaptive immune response. Activities are controlled by presence or absence of small cytoplasmic adapter proteins, SH2D1A/SAP and/or SH2D1B/EAT-2. Isoform 1 mediates NK cell activation through a SH2D1A-independent extracellular signal-regulated ERK-mediated pathway (PubMed:11698418). Positively regulates NK cell functions by a mechanism dependent on phosphorylated SH2D1B. Downstream signaling implicates PLCG1, PLCG2 and PI3K (PubMed:16339536). In addition to heterotypic NK cells-target cells interactions also homotypic interactions between NK cells may contribute to activation. However, in the absence of SH2D1B, inhibits NK cell function. Acts also inhibitory in T-cells (By similarity). May play a role in lymphocyte adhesion (PubMed:11802771). In LPS-activated monocytes negatively regulates production of proinflammatory cytokines (PubMed:23695528).[UniProtKB/Swiss-Prot Function]

**Product images:**


Circular map for RG236550