

Product datasheet for **RG237234**

PAX5 (NM_001280555) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PAX5 (NM_001280555) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	PAX5
Synonyms:	ALL3; BSAP
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG237234 representing NM_001280555. Blue=ORF Red=Cloning site Green=Tag(s)

```
GCTCGTTT TAGTGAACCGTCAGAATTTTGT AATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCC CGCATCGCC
ATGGATTTAGAGAAAAATTATCCGACTCCTCGACCAGCAGGACAGGACATGGAGGAGTGAATCAGCTT
GGGGGGTTTTTGTGAATGGACGGCCACTCCCGGATGTAGTCCGCCAGAGGATAGTGGAAC TTGCTCAT
CAAGGTGTCAGGCCCTGCGACATCTCCAGGCAGCTTCGGGT CAGCCATGGTTGTGCAGAAAATCTT
GGCAGGATCATCCGACAAAAGTACAGCAGCCACCCAACCAACCAGTCCCAGCTTCCAGTCACAGCATA
GTGTCCACTGGCTCCGTGACGCAGGTGTCTCGGTGAGCAGGATTCCGGCCGGCTCGTCTACTCCATC
AGCGGCATCCTGGGCATCACGTCCCCAGCGCCGACCAACAAGCGCAAGAGAGACGAAGTATTCAG
GAGTCTCCGGTGCCGAACGGCCACTCGCTTCCGGGCAGAGACTTCTCCGGAAGCAGATGCGGGGAGAC
TTGTTACACAGCAGCAGCTGGAGGTGCTGGACCGCGTGTGAGAGGCAGCACTACTCAGACATCTTC
ACCACCACAGAGCCCATCAAGCCCGAGCAGACCACAGAGTATTCAGCCATGGCCTCGCTGGTGGG
CTGGACGACATGAAGGCCAATCTGGCCAGCCCCACCCCTGCTGACATCGGGAGCAGTGTGCCAGGCCCG
CAGTCTACCCCATTTGTGACAGGGAGTGAGTTTTCCGGGAGTCCCTACAGCCACCCTCAGTATTCCTCG
TACAACGACTCCTGGAGGTTCCCAACCCGGGGCTGCTTGCTCCCCCTACTATTATAGCGCTGCCGCC
CGAGGAGCCGCCACCTGCAGCCGCCACTGCCTATGACCGTCAC
ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAAAC
```



[View online »](#)

Protein Sequence: >Peptide sequence encoded by RG237234
 Blue=ORF Red=Cloning site Green=Tag(s)

MDLEKNYPTPRTSRTGHGGVNQLGGVFNVRPLPDVVRQRIVELAHQGVRPCDISRQLRVSHGCVSKIL
 GRIIRTKVQQPPNPVPASSHSIVSTGSVTQVSSVSTDSAGSSYSISGILGITSPSADTNKRKRDEGIQ
 ESPVPNGHSLPGRDFLRKQMRGDLFTQQQLEVLDRVFERQHYSDIFTTTEPIKPEQTTEYSAMASLAGG
 LDDMKANLASPTPADIGSSVPGPQSYPIVTVGSEFSGSPYSHPQYSSYNDSWRFNPNGLLGSPYYYSAAA
 RGAAPPAATAYDRH
 TRTRPLEMESDESGLPAMEIECRITGTLNGVEFELVGGGEGTPEQGRMTNKMKSTKGALTFSPYLLSHV
 MGYGFYHFGTYPSTYENPFLHAINNGGYTNTRIEKYEDGGVLHVSFSYRYEAGRVIGDFKVMGTGFPEP
 SVIFTDKIIIRSNATVEHLHPMGDNDLDGSFTRTFSLRDGGYSSVVDSHMHFKSAIHPSILQNGGPMFA
 FRRVEEDHSNTELGIVEYQHAFKTPDADAGEERV

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001280555

ORF Size: 873 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

RefSeq: [NM_001280555.2](#)


RefSeq Size: 8609 bp

RefSeq ORF: 876 bp

Locus ID: 5079
UniProt ID: [Q02548](#)
Cytogenetics: 9p13.2
Protein Families: Transcription Factors
MW: 31.8 kDa

Gene Summary: This gene encodes a member of the paired box (PAX) family of transcription factors. The central feature of this gene family is a novel, highly conserved DNA-binding motif, known as the paired box. Paired box transcription factors are important regulators in early development, and alterations in the expression of their genes are thought to contribute to neoplastic transformation. This gene encodes the B-cell lineage specific activator protein that is expressed at early, but not late stages of B-cell differentiation. Its expression has also been detected in developing CNS and testis and so the encoded protein may also play a role in neural development and spermatogenesis. This gene is located at 9p13, which is involved in t(9;14)(p13;q32) translocations recurring in small lymphocytic lymphomas of the plasmacytoid subtype, and in derived large-cell lymphomas. This translocation brings the potent E-mu enhancer of the IgH gene into close proximity of the PAX5 promoter, suggesting that the deregulation of transcription of this gene contributes to the pathogenesis of these lymphomas. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2013]

Product images:



Circular map for RG237234