

Product datasheet for **RG237397**

CCR7 (NM_001301714) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	CCR7 (NM_001301714) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	CCR7
Synonyms:	BLR2; CC-CKR-7; CCR-7; CD197; CDw197; CMKBR7; EBI1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG237397 representing NM_001301714. Blue=ORF Red=Cloning site Green=Tag(s)

```
GCTCGTTTAGTGAACCGTCAGAATTTGTAAACGACTCACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGTACTCCATATTTGTTTCGTGGCCTACTGGGCAATGGGCTGGTTCGTGTTGACCTATATCTATTTT
AAGAGGCTCAAGACCATGACCGATACCTACCTGCTCAACCTGGCGGTGGCAGACATCCTCTTCTCCTG
ACCCTTCCCTTCTGGCCTACAGCGCGGCCAAGTCCTGGGTCTTCGGTGTCCACTTTTGAAGCTCATC
TTTGCCATCTACAAGATGAGCTTCTTCAGTGGCATGCTCCTACTTCTTTGCATCAGCATTGACCGCTAC
GTGGCCATCGTCCAGGCTGTCTCAGCTCACCGCCACCGTGCCTCCGCTCCTTCTCATCAGCAAGCTGTCC
TGTGTGGGCATCTGGATACTAGCCACAGTGTCTCCATCCAGAGCTCCTGTACAGTGACCTCCAGAGG
AGCAGCAGTGAGCAAGCGATGCGATGCTCTCTCATCACAGAGCATGTGGAGGCCTTTATCACCATCCAG
GTGGCCAGATGGTATCGGCTTTCTGGTCCCCCTGCTGGCCATGAGCTTCTGTTACCTTGTTCATCATC
CGCACCTGCTCCAGGCACGCAACTTTGAGCGCAACAAGGCCATCAAGGTGATCATCGTGTGGTTCGTG
GTCTTCATAGTCTTCCAGCTGCCCTACAATGGGGTGGTCTGGCCAGACGGTGGCCAACCTCAACATC
ACCAGTAGCACCTGTGAGCTCAGTAAGCAACTCAACATCGCCTACGACGTACCTACAGCCTGGCCTGC
GTCCGCTGCTGCGTCAACCCTTCTTGTACGCCTTCATCGGCGTCAAGTTCGGCAACGATCTCTTCAAG
CTCTTCAAGGACCTGGGCTGCCTCAGCCAGGAGCAGCTCCGGCAGTGGTCTTCTGTGCGCACATCCGG
CGCTCCTCCATGAGTGTGGAGGCCGAGACCACCACCTTCTCCCCA
ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAAAC
```



[View online »](#)

Protein Sequence: >Peptide sequence encoded by RG237397
 Blue=ORF Red=Cloning site Green=Tag(s)


MYSIICFVLLGNGLVVLTYIYFKRLKMTDITYLLNLAVADILFLLTLPFWAYSAAKSWVFGVHFCKLI
 FAIYKMSFFSGMLLLLCISIDRYVAIVQAVSAHRHRARVLLISKLSVGIWILATVLSIPELLYSDLQR
 SSSEQAMRCSLITEHVEAFITIQVAMVIGFLVPLLAMSFICYLVIIRTLQARNFERNKAIKVIIAVVV
 VFIVFQLPYNGVLAQTVANFNITSSTCELSKQLNIADVITYSLACVRCVNPFLYAFIGVKFRNDLFLK
 LFKDLGCLSQEQLRQWSSCRHIRRSMSEAEATTTTFSP
 TRTRPLEMESDESGLPAMEIECRITGTLNGVEFELVGGGEGTPEQGRMTNKMSTKGALTFSPYLLSHV
 MGYGFYHFGTYPSTYENPFLHAINNGGYNTRIEKYEDGGVLHVSFSYRYEAGRVIGDFKVMGTGFPEP
 SVIFTDKIIRSNATVEHLHPMGDNDLDGSFTRTFSLRDGGYSSVVDSHMHFKSAIHPSILQNGGPMFA
 FRRVEEDHSNTELGIVEYQHAFKTPDADAGEERV

Restriction Sites: SgfI-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_001301714

ORF Size:	945 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
RefSeq:	NM_001301714.2
RefSeq Size:	2157 bp
RefSeq ORF:	948 bp
Locus ID:	1236
UniProt ID:	P32248
Cytogenetics:	17q21.2
Protein Families:	Druggable Genome, GPCR, Transmembrane
Protein Pathways:	Chemokine signaling pathway, Cytokine-cytokine receptor interaction
MW:	36.2 kDa
Gene Summary:	The protein encoded by this gene is a member of the G protein-coupled receptor family. This receptor was identified as a gene induced by the Epstein-Barr virus (EBV), and is thought to be a mediator of EBV effects on B lymphocytes. This receptor is expressed in various lymphoid tissues and activates B and T lymphocytes. It has been shown to control the migration of memory T cells to inflamed tissues, as well as stimulate dendritic cell maturation. The chemokine (C-C motif) ligand 19 (CCL19/ECL) has been reported to be a specific ligand of this receptor. Signals mediated by this receptor regulate T cell homeostasis in lymph nodes, and may also function in the activation and polarization of T cells, and in chronic inflammation pathogenesis. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Sep 2014]