

## Product datasheet for **RN201287**

### **Rnps1 (NM\_001011890) Rat Untagged Clone**

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Rnps1 (NM\_001011890) Rat Untagged Clone  
**Tag:** Tag Free  
**Symbol:** Rnps1  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Fully Sequenced ORF:** >RN201287 representing NM\_001011890  
**Red**=Cloning site **Blue**=ORF **Orange**=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGATTTATCAGGAGTGAAAAAGAAGAGCTTGCTAGGAGTCAAAGAGAATAATAAAAAGTCCAGCACTA  
GGGCTCCTTCTCTACCAAACGAAAGGACCGCTCTGATGAGAAGTCCAAGGATCGATCTAAAGATAAAGG  
GACCACTAAAGAGTCCAGTGAGAAGGATCGTGGCAGAGATAAGACCCGGAAGAGACGCAGTGCTTCAAGC  
GGAAGCAGCAGCACCAGGTCCAGGTCCAGTCAACCTCCAGCTCGGGCTCCAGCACCAGCACGGGCTCGA  
GCAGTGGCTCTAGCTCGTCTTCTGCGTCCAGCCGCTCAGGAAGCTCCAGCACATCCCGGAGCTCCAGTTC  
CAGCAGCTCCTCCGGCTCCCCAAGCCCTTCTCGGCGCAGACATGACAACAGGCGGGCTTCCCGTCCAAA  
TCCAAACCACCTAAAAGAGATGAAAAAGAGAGGAAAAGGCGGAGCCCTTACCTAAACCCACCAAAGTGC  
ACATTGGGAGGCTCACCAGGAATGTGACCAAGGATCATATCATGGAAATATTTTCTACTTACGGGAAAAT  
TAAAATGATTGACATGCCTGTGGAAAGGATGCATCCTCATCTATCCAAAGGCTATGCGTACGTGGAATTC  
GAGAATCCGGATGAAGCAGAGAAGGCACTGAAACACATGGATGGAGGACAAATAGATGGCCAAGAGATCA  
CTGCTACTGCGGTGTTGGCACCCTGGCCTCGTCCACCGCCTCGGCGATTACGCCACCCAGGAGAATGCT  
TCCACCGCCTCCCATGTGGCGTAGTCAACCCACGGATGAGGAGAAGGTCTCGATCCCCAAGACGCAGG  
TCCCCTGTGCGAAGGAGGTCTCGATCGCCTGGCCGCCCGCCACAGGAGCCGATCCAGCTCCAACCTCT  
CCCGAT**TAA**

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** Sgfl-MluI  
**ACCN:** NM\_001011890  
**Insert Size:** 918 bp



<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u><a href="#">NM_001011890.1</a></u> , <u><a href="#">NP_001011890.1</a></u>
<b>RefSeq Size:</b>	1853 bp
<b>RefSeq ORF:</b>	918 bp
<b>Locus ID:</b>	287113
<b>UniProt ID:</b>	<u><a href="#">Q6AYK1</a></u>
<b>Cytogenetics:</b>	10q12
<b>Gene Summary:</b>	<p>Part of pre- and post-splicing multiprotein mRNP complexes. Auxiliary component of the splicing-dependent multiprotein exon junction complex (EJC) deposited at splice junction on mRNAs. The EJC is a dynamic structure consisting of core proteins and several peripheral nuclear and cytoplasmic associated factors that join the complex only transiently either during EJC assembly or during subsequent mRNA metabolism. Component of the ASAP and PSAP complexes which bind RNA in a sequence-independent manner and are proposed to be recruited to the EJC prior to or during the splicing process and to regulate specific excision of introns in specific transcription subsets. The ASAP complex can inhibit RNA processing during in vitro splicing reactions. The ASAP complex promotes apoptosis and is disassembled after induction of apoptosis. Enhances the formation of the ATP-dependent A complex of the spliceosome. Involved in both constitutive splicing and, in association with SRP54 and TRA2B/SFRS10, in distinctive modulation of alternative splicing in a substrate-dependent manner. Involved in the splicing modulation of BCL2L1/Bcl-X (and probably other apoptotic genes); specifically inhibits formation of proapoptotic isoforms such as Bcl-X(S); the activity is different from the established EJC assembly and function. Participates in mRNA 3'-end cleavage. Involved in UPF2-dependent nonsense-mediated decay (NMD) of mRNAs containing premature stop codons. Also mediates increase of mRNA abundance and translational efficiency. Binds spliced mRNA 20-25 nt upstream of exon-exon junctions (By similarity). [UniProtKB/Swiss-Prot Function]</p>