

Product datasheet for RN203375

Nudt2 (NM_207596) Rat Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Nudt2 (NM_207596) Rat Untagged Clone

Tag: Tag Free Symbol: Nudt2

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >RN203375 representing NM_207596

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul ACCN: NM 207596

Insert Size: 444 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 207596.1</u>, <u>NP 997479.1</u>

RefSeq Size: 939 bp
RefSeq ORF: 444 bp
Locus ID: 297998
UniProt ID: Q6PEC0
Cytogenetics: 5q22

Gene Summary: member of the nudix family of hydrolase proteins, and of the subfamily of nucleotide

pyrophosphatases containing the MutT sequence motif; hydrolyzes Ap4a to yield AMP and

ATP [RGD, Feb 2006]