

Product datasheet for **RN204917**

Pnrc2 (NM_001103360) Rat Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Pnrc2 (NM_001103360) Rat Untagged Clone
Tag: Tag Free
Symbol: Pnrc2
Synonyms: MGC93828
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >RN204917 representing NM_001103360
Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGGGTGGTGGAGAGAGGTATAACATTCCAGACCCTCAATCTAGAAATGCTAGCAAGAACCAACAACAGC
ACAATAGACAGAAGACCAAGGATCAGAATCCCAGATGAAGATCGTTCATAAGAAAAAGGAAAGAGGACA
TGGGTACAATCCATCGGCAGTGC AAAATGGGGAAAAACCAAGAGCCTTCCAACAACCTCAACTGGAAT
GCTAGCTTATCAAGTCTAGCTTCTTTTAAAGTCTCAAGCTAGTCAGAACTATGCTGGAGCCAAATTTA
GTGAACCAACCATACCAAGTGTCTCCCAAGCCACCAAGCCACTGGGTTTCATGTTTCCTTGAACCTTC
AGATAAGGAAACGATGACATTTCAACTTAAACCTTACTTAAAGTACAGGTAA

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI
ACCN: NM_001103360
Insert Size: 405 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001103360.1](#), [NP_001096830.1](#)

RefSeq Size: 1899 bp

RefSeq ORF: 405 bp

Locus ID: 100125373

UniProt ID: [Q66HE1](#)

Cytogenetics: 5q36

Gene Summary: Involved in nonsense-mediated mRNA decay (NMD) by acting as a bridge between the mRNA decapping complex and the NMD machinery. May act by targeting the NMD machinery to the P-body and recruiting the decapping machinery to aberrant mRNAs. Required for UPF1/RENT1 localization to the P-body. Plays a role in glucocorticoid receptor-mediated mRNA degradation by interacting with the glucocorticoid receptor NR3C1 in a ligand-dependent manner when it is bound to the 5' UTR of target mRNAs and recruiting the RNA helicase UPF1 and the mRNA-decapping enzyme DCP1A, leading to RNA decay. Also acts as a nuclear receptor coactivator. May play a role in controlling the energy balance between energy storage and energy expenditure.[UniProtKB/Swiss-Prot Function]