

## **Product datasheet for RN206657**

## Mafg (NM\_022386) Rat Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Mafg (NM 022386) Rat Untagged Clone

Tag: Tag Free
Symbol: Mafg

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >RN206657 representing NM\_022386

489 bp

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** Sgfl-Mlul NM\_022386

Insert Size:

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 022386.2, NP 071781.1

RefSeq Size: 1638 bp RefSeq ORF: 489 bp Locus ID: 64188 **UniProt ID:** Q76MX4 Cytogenetics: 10q32.3

**Gene Summary:** increased expression of splice variant MafG-2 is detected in response to decreased

extracellular pH; may play a role in signal transduction of extracellular protons [RGD, Feb

2006]