

Product datasheet for RN206748

Slc4a8 (NM_199497) Rat Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Slc4a8 (NM_199497) Rat Untagged Clone

Tag: Tag Free Symbol: Slc4a8

Mammalian Cell Neomycin

Selection:

Vector: pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

Restriction Sites: Sgfl-Mlul ACCN: NM_199497

Insert Size: 3204 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 199497.2</u>, <u>NP 955791.1</u>

 RefSeq Size:
 3626 bp

 RefSeq ORF:
 3204 bp

 Locus ID:
 315311



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Slc4a8 (NM_199497) Rat Untagged Clone - RN206748

UniProt ID: Q6RVG2

Cytogenetics: 7q36

Gene Summary: may act as a bicarbonate transporter [RGD, Feb 2006]

Transcript Variant: This variant (1) represents the longest transcript and encodes the longest isoform (1). Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The

genomic coordinates used for the transcript record were based on alignments.