

Product datasheet for **RN209750**

Nrep (NM_178096) Rat Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Nrep (NM_178096) Rat Untagged Clone
Tag: Tag Free
Symbol: Nrep
Synonyms: P311
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >RN209750 representing NM_178096
Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGTTTACTACCGGAACCTCTGGTCTGGTCTAGCCAGGAGCCATTTCCATACAAGGAAATGGAGGGAG
GTCTTACTAAGGGAAGACTTCTGTGCCTAAGGAAGTGAACCGCAAGAAGATGGATGAGCCCATCGCTGC
CTCGCTGATTCCGCCGGCAACCATGAACCCGGCTCCCCAGCTATCGGTTACCTCCACCCTTTTAA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI
ACCN: NM_178096
Insert Size: 207 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_178096.3](#), [NP_835197.1](#)

RefSeq Size: 435 bp

RefSeq ORF: 207 bp

Locus ID: 338475

UniProt ID: [Q80Z34](#)

Cytogenetics: 18p12

Gene Summary: may facilitate nerve regeneration [RGD, Feb 2006]