

## Product datasheet for RN209820

### Med22 (NM\_001077679) Rat Untagged Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Med22 (NM\_001077679) Rat Untagged Clone  
**Tag:** Tag Free  
**Symbol:** Med22  
**Synonyms:** RGD1564893; Surf5  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Fully Sequenced ORF:** >RN209820 representing NM\_001077679  
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGCC**

ATGGCCCAGCAGAGAGCCCTGCCCCAGAGCAAAGAGACGCTACTGCAGTCATATAACAAGCGGCTCAAAG  
 ACGACATCAAGTCCATCATGGACAACCTCACCGAGATCATCAAACGGCCAAGATTGAGGATGAGACGCA  
 GGTGTCGAGGGCTACTCAGGGCGAACAGGACAACACGAGATGCACGTTTCGAGCTGCCAACATTGTCGG  
 GCTGGTGAGTCGCTGATGAAGCTGGTATCTGATCTCAAGCAGTTTCTGATCCTCAATGACTTCCCATCTG  
 TGAATGAAGCCATTGACCAGCGTAACCAGCAGCTTCGAGCCCTGCAGGAGGAGTGTGACCGCAAGCTCAT  
 CACCCTGCGGGATGAGGTCTCCATTGACCTGTATGAGCTGGAGGAGGAGTACTACTCGTCCAGCTCAAGT  
 CTCTGCGAAGCTAATGACCTGCCTCTATGTGAAGCTTACTGGAGGCTGGACCTCGACACAGATTCTGCTG  
 ATGGCCTCTCAGCCCCTTTCTGGCGTCCCCGGAGACTGGTGTGGTCCCTTGCAGTCTGCAGCCCTGT  
 CCACTCCCATGGTGGTGGCCCTGGCCCTACAGAACACACCT**GTA**

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** Sgfl-Mlul  
**ACCN:** NM\_001077679  
**Insert Size:** 603 bp



[View online »](#)

<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u><a href="#">NM_001077679.1</a></u> , <u><a href="#">NP_001071147.1</a></u>
<b>RefSeq Size:</b>	1415 bp
<b>RefSeq ORF:</b>	603 bp
<b>Locus ID:</b>	499762
<b>UniProt ID:</b>	<u><a href="#">A0JPN6</a></u>
<b>Cytogenetics:</b>	3p13
<b>Gene Summary:</b>	Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors (By similarity). [UniProtKB/Swiss-Prot Function]