

Product datasheet for RN210410

Babam1 (NM 001006964) Rat Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Babam1 (NM_001006964) Rat Untagged Clone

Tag: Tag Free Babam1 Symbol: Merit40 Synonyms: **Mammalian Cell**

Selection:

Neomycin

Vector: pCMV6-Entry (PS100001) E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >RN210410 representing NM_001006964

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

CGATTTCAGAGCCCAGGCCTCACACTCGCTCCAATCCTGAGGGGGCTGAGGACCGGGCCATCGGGGCTCA GGCCAATGTGGGTAGCCGCAGCGAGGGCGAAGGTGAGGCAGCCACTGCAGATGATGGGGCGGCCAGTGTC CCGGGAGCTGTGCCCAAGCCCTGGCAGGTACCAGCACCAGCATCTGAGGTCCAGATTCGAACACCAAGGG TCAACTGTCCAGAGAAAGTGATCATCTGTCTGGATCTTTCAGAGGAGATGTCTGTGCCAAAGCTGGAGTC CTTTAATGGGTCCAGAACAACGCCCTGAATGTGTCTCAGAAGATGGTTGAGATGTTTGTGCGCACGAAG CACAAGATTGACAAGAGCCACGAGTTTGCCTTGGTCGTAGTGAACGACGACTCTGCCTGGTTGTCTGGCC TGACCTCTGACCCACGTGAACTCTGCAGCTGCCTGTACGACCTAGAGACGGCATCCTGCTCCACATTCAA TTTGGAAGGCCTCTTCAGCCTCATCCAGCAGAAGACTGAGCTGCCAGTCACAGAGAATGTGCAAACCATC CCACCCCCTACGTCGTGCGCACCATCCTGGTCTACAGCCGCCCACCCTGCCAGCCCCAGTTCTCCTTGA CTGAGCCCATGAAGAAGATGTTCCAATGTCCCTACTTCTTCTTCGACATCATTTACATCCACAGTGGCCC TGAGGAAAAGGAAGACGATATGAGCTGGAAGGACATGTTCGCCTTCATGGGCAGTCTGGACACCAAGGGC ACCAGCTACAAGTATGCAGTAGCACTTGCTGGCCCCGCCCTGGAGCTGCACAACTGCGTGGCCAAGTTGC TGGCCCACCCGCTGCAGAGGCCCTGCCAGAGCCACGCGAGCTATAGCCTGCTGGAAGAGGACGAAGAGGC CGGTGAGGGGGAGGCCACTGTGTGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

Babam1 (NM_001006964) Rat Untagged Clone - RN210410

ACCN: NM_001006964

Insert Size: 1005 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001006964.1</u>, <u>NP 001006965.1</u>

 RefSeq Size:
 1368 bp

 RefSeq ORF:
 1005 bp

 Locus ID:
 290631

 UniProt ID:
 Q5XII6

 Cytogenetics:
 16p14

Gene Summary: Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked

ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it is required for the complex integrity and its localization at DSBs. Component of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates. In these 2 complexes, it is probably required to maintain the stability of BABAM2 and help the 'Lys-63'-linked deubiquitinase activity mediated by BRCC3/BRCC36 component. The BRISC complex is required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitinating NUMA1. Plays a role in interferon signaling via its role in the

deubiquitination of the interferon receptor IFNAR1; deubiquitination increases IFNAR1 activity by enhancing its stability and cell surface expression. Down-regulates the response to bacterial lipopolysaccharide (LPS) via its role in IFNAR1 deubiquitination. [UniProtKB/Swiss-

Prot Function]