

Product datasheet for RN211262

Hopx (NM_133621) Rat Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Hopx (NM_133621) Rat Untagged Clone
Tag: Tag Free
Symbol: Hopx
Synonyms: Gllg15b; Hod; Obl
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >RN211262 representing NM_133621
Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCCGCGATCGCC

ATGTCGGCGCAGACTGCGAGCGGCCCCACGGAGGACCAGGTGGAGATCCTGGAGTACAACCTCAACAAGG
TCAACAAGCACCCGACCCACCCACGCTGTGCCTCATCGACCCGAGGCGGGCCTCACGGAGGAGCAGAC
GCAGAAATGGTTAAGCAGCGCCTGGCGGAGTGGCGGCGGTGAGAAGGCCTGCCTTCGGAATGCAGATCG
GTCACGGACTAG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-MluI
ACCN: NM_133621
Insert Size: 222 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_133621.2](#), [NP_598305.2](#)

RefSeq Size: 1065 bp

RefSeq ORF: 222 bp

Locus ID: 171160

UniProt ID: [Q78ZR5](#)

Cytogenetics: 14p11

Gene Summary: Atypical homeodomain protein which does not bind DNA and is required to modulate cardiac growth and development. Acts via its interaction with SRF, thereby modulating the expression of SRF-dependent cardiac-specific genes and cardiac development. Prevents SRF-dependent transcription either by inhibiting SRF binding to DNA or by recruiting histone deacetylase (HDAC) proteins that prevent transcription by SRF. Overexpression causes cardiac hypertrophy. Acts as a co-chaperone for HSPA1A and HSPA1B chaperone proteins and assists in chaperone-mediated protein refolding.[UniProtKB/Swiss-Prot Function]