

Product datasheet for **RN211903**

Serp1 (NM_030835) Rat Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Serp1 (NM_030835) Rat Untagged Clone
Tag: Tag Free
Symbol: Serp1
Synonyms: RAMP4
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >RN211903 representing NM_030835
Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGTCGCCAAGCAGAGGATCCGTATGGCCAACGAGAAGCACAGCAAGAACATAACTCAGCGGGCAACG
TCGCTAAGACCTCGAAAAATGCCCCGAAGAAAAGGCGTCGGTAGGACCCTGGTTATTGGCCCTCTTCAT
TTTTGTCGTTTGGATCTGCAATTTCCAGATTATCAAGATCAGGATGGGCATGTA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI
ACCN: NM_030835
Insert Size: 201 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_030835.2](#), [NP_110462.1](#)

RefSeq Size: 2442 bp

RefSeq ORF: 201 bp

Locus ID: 80881

UniProt ID: [Q9R2C1](#)

Cytogenetics: 2q26

Gene Summary: may be involved in glycosylation modification of secretory and membrane proteins including glycosylation of MHC class II-associated invariant chain (Ii) [RGD, Feb 2006]