

Product datasheet for **RN213619**

Mapkap1 (NM_001011964) Rat Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Mapkap1 (NM_001011964) Rat Untagged Clone
Tag:	Tag Free
Symbol:	Mapkap1
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

Fully Sequenced ORF: >RN213619 representing NM_001011964
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGCCTTCTGGACAATCCAATCATCTAGCTCATATTCGACAGTCACATGTGACCAGTGATGACA
 CGGGAATGTGAGATGGTTCTCATTGATCATGACGTTGACCTAGAGAAGACTCATCCTCCGTCAGTGCC
 TGGAGACAGTGGGTCAGAAGTTCAGGGAAGCAGTGGTGAGACGCAGGGCTACATATACGCCCACTGTGT
 GATATTACCTCGAGCTGGGACTTTGGTATTAGAAGACGCTCAAACACAGCTCAAAGATTAGAACGACTCC
 GCAAAGAGAGACAAAACAGATCAAATGCAAAAATATTAGTGGAAAGAAAGAAATCTAAACAATCAGC
 CCAGGAGTTAAAGTCACTGTTTGAATAAATCCCTCAAAGAGAAACCTCCAAGTTCGGGCAAGCAGTCC
 ATATTGTCTGTGCGCCTGGAACAGTGCCTCTGCAGCTGAATAACCCCTTAATGAGTACTCCAAGTTG
 ATGGCAAGGGTCACTGTGGTACAACGCAAGAAGATCGACGTCTACCTCCCCTGCACTCAAGCCA
 GGACAGACTGCTCCAATGACTGTGGTACCATGGCCAGCGCCAGGGTGCAGGACCTCATCGGGCTCATC
 TGTGGCAGTACACGAGTGAAGGACGGGAGCCAAAGCTCAATGACAACGTCAGTGCCTACTGCCTGCATA
 TTGCTGAGGATGATGGGGAGGTGGACACGGATTTCCACCCTGGATTCCAATGAACCCATTCAAGTT
 TGGCTTCACTACTTTGGCCCTGGTTGAAAAGTACTCTTCCCTGGTCTGACCTCCAAAGAGTCGCTCTTT
 GTTCGAATAAATGCTGCCATGGGTTCTCCCTCATCCAGTGGACAACACGAAGGTCACCATGAAGGAGA
 TCTTACTCAAGGCACTGAAACGAAGAAAAGGGTCCCAGAAAATTCAGGCCCTCAGTACCGCCTGGAGAA
 GCAGAGCGAGCCTAACATCGGTGTGGACCTGGAGAGCACGCTGGAGAGCCAGAAATGCCTGGGAGTTCTGC
 CTGGTTCGAGAGAACAGTTCAAGGGCAGACGGAGTTTTTCGAGGAGGATTACAAAATTGACATTGCTACAG
 TGCAGGATATGCTTAGCAGCCACCATTATAAGTCATTCAAAGTCAGCATGATCCACAGACTGCGGTTTAC
 AACTGACGTGCAGTTAGGTATCTCTGGAGACAAAGTGGAGATAGACCCTGTTACGAATCAGAAAGCCAGC
 ACTAAGTTTTGGATTAAGCAGAAGCCATCTCAATCGATTGTGACCTGCTCTGTGCTGTGACCTTGCCG
 AGGAGAAGAGCCCACTCACGGGCTTTAAGCTCACGTATCTAAGCAGTCACGACTATAAGCATCTCTA
 CTTGAGTCCGACGCAGTACCCTCAGTAAATCGTGCTCAAGGTTAACTACATCTGGAGTACAGAGCA
 AGCACTGCCCGTGTGACTATTTTGCTCAAAAACAAGAAAAGTGAACAGGCGGACGAGCTTCAGCTTCC
 AGAAGGAGAAGAAATCAGGGCAGCAGTGA

ACGGTACGGCGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI

ACCN: NM_001011964

Insert Size: 1569 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001011964.1](#), [NP_001011964.1](#)

RefSeq Size: 2400 bp

RefSeq ORF: 1569 bp

Locus ID: 296648

UniProt ID: [Q6AYF1](#)

Cytogenetics: 3p11

Gene Summary: Subunit of mTORC2, which regulates cell growth and survival in response to hormonal signals. mTORC2 is activated by growth factors, but, in contrast to mTORC1, seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at 'Ser-421'. mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657'. Within mTORC2, MAPKAP1 is required for complex formation and mTORC2 kinase activity. MAPKAP1 inhibits MAP3K2 by preventing its dimerization and autophosphorylation. Inhibits HRAS and KRAS signaling. Enhances osmotic stress-induced phosphorylation of ATF2 and ATF2-mediated transcription. Involved in ciliogenesis, regulates cilia length through its interaction with CCDC28B independently of mTORC2 complex (By similarity).[UniProtKB/Swiss-Prot Function]