

Product datasheet for RN215718

Med20 (NM_001277367) Rat Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Med20 (NM_001277367) Rat Untagged Clone
Tag:	Tag Free
Symbol:	Med20
Synonyms:	Trfp; Usp49
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>RN215718 representing NM_001277367 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGGAGTGACGTGTGTGCCAGATGCCTGTGGCTGAGGGCAAGAGTGTCCAGCAGACAGTGGAGCTCC
TCACCAAGAAATTGGAGATGCTTGGAGCAGAGAAACAAGGAACATTTTGTGTGGACTGTGAACTTACCA
CACAGCTGCCTCCACCCTGGGCAGCCAAGGTGGAGTATGGCCCTGTGTGGTAGC**TAG**

ACGGTACGGCGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites:	Sgfl-MluI
ACCN:	NM_001277367
Insert Size:	198 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.



[View online »](#)

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001277367.1</u> , <u>NP_001264296.1</u>
RefSeq Size:	1710 bp
RefSeq ORF:	198 bp
Locus ID:	316209
Cytogenetics:	9q12
Gene Summary:	<p>Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors (By similarity). [UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (2) lacks an alternate exon in the coding region, which results in a frameshift, compared to variant 1. It encodes isoform 2, which has a shorter and distinct C-terminus, compared to isoform 1. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>