

## Product datasheet for **RN216115**

### Cidec (NM\_001244798) Rat Untagged Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Cidec (NM\_001244798) Rat Untagged Clone  
**Tag:** Tag Free  
**Symbol:** Cidec  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Cell Selection:** Neomycin  
**Fully Sequenced ORF:** >RN216115 representing NM\_001244798  
Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGC**C

ATGGACTATGCTATGAAGTCTCTCAGCCTTCTCTACCCTAGGTCAGTGTCCAGGCATGTAGCAGTGAGCA  
CTGCAGTGGTGACCCAACAGCTGGTGTCTGAGCCAGCCGGGAGACCCCCAGGGCAAGACCCTGTCGTGT  
TAGCACCGCAGATCGGAAGGTTTCGAAAGGCATCATGGCCACAGCTTGGAGGACCTCCTGGCAAGGTC  
CAAGACATCTTGAAGCTTAAAGACAAGCCCTTCTCCCTGGTGTGGAGGAAGATGGCACAATCGTGGAGA  
CAGAAGAATACTTCCAAGCCCTACCAAGAGATACAGTGTTCATGGTCTGCAGAAGGGGCAGAAGTGGAA  
GTCCCCATCAGAACAGCGCAAGAAGAAAGCCAGCTATCCCTTCCAGAAAGCCAACTAAGAAGATCGAC  
GTGGCCCGGGTAACCTTTGACCTGTACAAGCTGAACCCTCAGGACTTCATCGGCTGCCTGAACGTGAAGG  
CAACCCTCTATGACACATACTCGCTTTCATATGACCTGCACTGCTACAGGGCCAAACGCATCGTGAAGGA  
AATGCTCCGCTGGACTCTCTTCAATGCAAGCCACAGGCCACATGCTGCTTGGCACCTCCAGTTACATG  
CAGCAGTTCCTGGATGCCACTGAGGAAGAAGCCCTCCAAGGCCAAGGCCTCCCTCCTCCCTGCCTGTC  
TGAAGATGCTGCAATGA

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** Sgfl-Mlul  
**ACCN:** NM\_001244798  
**Insert Size:** 717 bp



[View online »](#)

<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>OTI Annotation:</b>	Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u><a href="#">NM_001244798.1</a></u> , <u><a href="#">NP_001231727.1</a></u>
<b>RefSeq Size:</b>	1655 bp
<b>RefSeq ORF:</b>	717 bp
<b>Locus ID:</b>	500292
<b>UniProt ID:</b>	<u><a href="#">Q5XI33</a></u>
<b>Cytogenetics:</b>	4q42
<b>Gene Summary:</b>	<p>Binds to lipid droplets and regulates their enlargement, thereby restricting lipolysis and favoring storage. At focal contact sites between lipid droplets, promotes directional net neutral lipid transfer from the smaller to larger lipid droplets. The transfer direction may be driven by the internal pressure difference between the contacting lipid droplet pair. Its role in neutral lipid transfer and lipid droplet enlargement is activated by the interaction with PLIN1. May act as a CEBPB coactivator in the white adipose tissue to control the expression of a subset of CEBPB downstream target genes, including SOCS1, SOCS3, TGFB1, TGFBR1, ID2 and XDH. When overexpressed in preadipocytes, induces apoptosis or increases cell susceptibility to apoptosis induced by serum deprivation or TGFB treatment. The physiological significance of its role in apoptosis is unclear (By similarity). May play a role in the modulation of the response to osmotic stress by preventing NFAT5 to translocate into the nucleus and activate its target genes expression.[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (3) differs in the 5' exon, which results in a downstream AUG start codon, compared to variant 1. The resulting isoform (2) is shorter at the N-terminus than isoform 1.</p>