

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RR200006L4V

## Ost4 (NM\_001134690) Rat Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Ost4 (NM_001134690) Rat Tagged ORF Clone Lentiviral Particle
Symbol:	Ost4
Synonyms:	Ost4
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001134690
ORF Size:	111 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RR200006).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001134690.1, NP 001128162.1</u>
RefSeq Size:	570 bp
RefSeq ORF:	114 bp
Locus ID:	100188932
UniProt ID:	BOBLSO
Cytogenetics:	6q14



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Ost4 (NM_001134690) Rat Tagged ORF Clone Lentiviral Particle – RR200006L4V
Gene Summary:	Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial transfer of a defined glycan (Glc(3)Man(9)GlcNAc(2) in eukaryotes) from the lipid carrier dolichol- pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent
	polypeptide chains, the first step in protein N-glycosylation. N-glycosylation occurs
	cotranslationally and the complex associates with the Sec61 complex at the channel-forming

translocon complex that mediates protein translocation across the endoplasmic reticulum

(ER). All subunits are required for a maximal enzyme activity. Specifically involved in maintaining stability of STT3A-containing OST complexes.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US