



Product datasheet for RR200141L4

Nit1 (NM_182668) Rat Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Nit1 (NM_182668) Rat Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Nit1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RR200141).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.

ACCN: NM_182668


ORF Size: 981 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_182668.2 , NP_872609.2
RefSeq Size:	1391 bp
RefSeq ORF:	984 bp
Locus ID:	289222
UniProt ID:	Q7TQ94
Cytogenetics:	13q24
Gene Summary:	The protein encoded by this gene is a member of the nitrilase enzyme superfamily. Nitrilases are nonpeptidic C-N hydrolases. In mammals, nitrilases form small molecule C-N bonds such as acid amides, carbamates and ureas. Nit1-deficient mice display phenotypes including increased cell proliferation, resistance to DNA damage, and an increased number of N-nitrosomethylbenzylamine-induced murine forestomach tumors, suggesting a tumor suppressor function. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2015]

Product images:



Circular map for RR200141L4