

Product datasheet for **RR200392**

Gpx4 (NM_001039849) Rat Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Gpx4 (NM_001039849) Rat Tagged ORF Clone
Symbol:	Gpx4
Synonyms:	gpx-4; Gshpx-4; Phgpx; snGpx
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RR200392 representing NM_001039849 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGGCCGCGCGCCGCCCGCAAGCGGGGACGCTGCAGACAGCGCGCCGGTCCCCGGGAGGCCGGCGAC
GGCGTGAACCTGGACGCCAAAGTCCTAGGAAGCGCCAGGCCCTCGGAGGAGGAGAGCTCGCGCGCGCCG
CCGAGGAGGGCGCGCCCTCGCCGGATGGAGCCATTCCCGAGCCTTTCAACCCGCGCCCTCTGCTGCAG
GACCTTCCCAGACCAGCAACAGCCACGAGTTCCTGGGCTTGTGTGCATCCCGCATGATTGGCGCTGTG
CGCGCTCCATGCACGAATTCGACGCCAAGGACATCGATGGGCACATGGTTGCCTGGATAAGTACAGGGG
TTGCGTGTGCATCGTCACCAACGTGGCCTCGCAATGAGGCAAAACCGACGTAAACTACACTCAGCTAGTC
GATCTGCATGCCCGATACGCCGAGTGTGGTTTACGAATCCTGGCCTTCCCTTGAACCAAGTTCGGGAGGC
AGGAGCCAGGAAGTAATCAAGAAATCAAGGAGTTTGCAGCCGGCTACAATGTCAGGTTTGACATGTACAG
CAAGATCTGTGTAATGGGGACGATGCCACCCACTGTGGAAATGGATGAAAGTCCAGCCCAAGGGCAGG
GGCATGCTGGGAAATGCCATCAAATGGAACCTTACCAAGTTTCTCATTGATAAGAACGGCTCGTGGTGA
AGCGCTATGGTCCATGGAGGAGCCCCAGGTGATAGAGAAGGACCTGCCGTGCTATCTC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)


Protein Sequence: >RR200392 representing NM_001039849
 Red=Cloning site Green=Tags(s)

MGRAAARKRGRCRQRGRSPGRRRRREPGRQSPRKRPGPRRRRARARRRRRARPRRMEPIPEPFNPRPLLQ
 DLPQTSNSHEFLGLCASRDDWRCARSMHEFAAKDIDGHMVCLDKYRGCVCIVTNVASQ*GKTDVNYTQLV
 DLHARYAECGLRILAFPCNQFGRQEPGSNQEIKEFAAGYNVRFDMYSKICVNGDDAHLWKWMKVQPKGR
 GMLGNAIKWNFTKFLIDKNGCVVKRYGPMEEPQVIEKDLPCYL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001039849

ORF Size: 759 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#) The expression of this clone is not guaranteed due to the nature of selenoproteins.

OTI Annotation: This clone encodes a selenoprotein containing the rare amino acid selenocysteine (Sec). Sec is encoded by UGA codon, which normally signals translational termination. Expression of this clone is not guaranteed due to the nature of selenoproteins.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001039849.3](#)

RefSeq Size: 1037 bp

RefSeq ORF: 762 bp

Locus ID: 29328


UniProt ID: [P36970](#)

Cytogenetics: 7q11

MW: 29.3 kDa

Gene Summary: The protein encoded by this gene belongs to the glutathione peroxidase family, members of which catalyze the reduction of hydrogen peroxide, organic hydroperoxides and lipid hydroperoxides, and thereby protect cells against oxidative damage. Several isozymes of this gene family exist in vertebrates, which vary in cellular location and substrate specificity. This isozyme has a high preference for lipid hydroperoxides and protects cells against membrane lipid peroxidation and cell death. It is also required for normal sperm development; thus, it has been identified as a 'moonlighting' protein because of its ability to serve dual functions as a peroxidase, as well as a structural protein in mature spermatozoa. Disruption of this gene in mouse spermatocytes is associated with male infertility. This isozyme is also a selenoprotein, containing the rare amino acid selenocysteine (Sec) at its active site. Sec is encoded by the UGA codon, which normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, designated the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon, rather than as a stop signal. Transcript variants resulting from alternative splicing or use of alternate promoters have been described to encode isoforms with different subcellular localization. Pseudogenes of this locus have been identified on chromosomes 1, 10 and 14. [provided by RefSeq, Jan 2019]

Product images:



Circular map for RR200392