

Product datasheet for RR200876L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Apln (NM_031612) Rat Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Apln (NM_031612) Rat Tagged ORF Clone Lentiviral Particle

Symbol: Apln Synonyms: Apel

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_031612

ORF Size: 231 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RR200876).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 031612.3, NP 113800.1

RefSeq Size: 3114 bp
RefSeq ORF: 234 bp
Locus ID: 58812
UniProt ID: Q9R0R3
Cytogenetics: Xq36







Gene Summary:

Endogenous ligand for the apelin receptor (APLNR) (PubMed:11336787, PubMed:11359874, PubMed:26611206). Drives internalization of the apelin receptor (PubMed:11359874). Apelin-36 dissociates more hardly than (pyroglu)apelin-13 from APLNR (PubMed:11336787). Hormone involved in the regulation of cardiac precursor cell movements during gastrulation and heart morphogenesis (By similarity). Has an inhibitory effect on cytokine production in response to T-cell receptor/CD3 cross-linking; the oral intake of apelin in the colostrum and the milk might therefore modulate immune responses in neonates (By similarity). Plays a role in early coronary blood vessels formation (By similarity). Mediates myocardial contractility in an ERK1/2-dependent manner (PubMed:26611206). May also have a role in the central control of body fluid homeostasis by influencing vasopressin release and drinking behavior (PubMed:10617103, PubMed:11359874).[UniProtKB/Swiss-Prot Function]