

## Product datasheet for RR201213L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Mtmr6 (NM\_001107268) Rat Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Mtmr6 (NM\_001107268) Rat Tagged ORF Clone Lentiviral Particle

Symbol: Mtmr6

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_001107268

ORF Size: 981 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RR201213).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 001107268.1</u>, <u>NP 001100738.1</u>

 RefSeq Size:
 3779 bp

 RefSeq ORF:
 984 bp

 Locus ID:
 305935

 Cytogenetics:
 15p12







## **Gene Summary:**

Phosphatase that acts on lipids with a phosphoinositol headgroup. Dephosphorylates phosphatidylinositol 3-phosphate (PtdIns(3)P) and phosphatidylinositol 3,5-bisphosphate. Binds with high affinity to phosphatidylinositol 3,5-bisphosphate (PtdIns(3,5)P2) but also to phosphatidylinositol 3-phosphate (PtdIns(3)P), phosphatidylinositol 4-phosphate (PtdIns(4)P), and phosphatidylinositol 5-phosphate (PtdIns(5)P), phosphatidic acid and phosphatidylserine (By similarity). Negatively regulates ER-Golgi protein transport (PubMed:23188820). Probably in association with MTMR9, plays a role in the late stages of macropinocytosis by dephosphorylating phosphatidylinositol 3-phosphate in membrane ruffles. Acts as a negative regulator of KCNN4/KCa3.1 channel activity in CD4(+) T-cells possibly by decreasing intracellular levels of phosphatidylinositol 3-phosphate. Negatively regulates proliferation of reactivated CD4(+) T-cells. In complex with MTMR9, negatively regulates DNA damage-induced apoptosis. The formation of the MTMR6-MTMR9 complex stabilizes both MTMR6 and MTMR9 protein levels (By similarity).[UniProtKB/Swiss-Prot Function]