

Product datasheet for RR201425L4

Akr1c2 (NM_001013057) Rat Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Akr1c2 (NM_001013057) Rat Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Akr1c2
Synonyms:	Akr1c21
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RR201425).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	


Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.



[View online »](#)

Plasmid Map:


ACCN: NM_001013057

ORF Size: 954 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001013057.1](#)

RefSeq Size: 1398 bp

RefSeq ORF: 957 bp

Locus ID: 291283

UniProt ID: [Q6AYQ2](#)

Cytogenetics: 17q12.2

Gene Summary: NADP-dependent 17-alpha-hydroxysteroid dehydrogenase that converts 5-alpha-androstane-3,17-dione into androsterone. Has lower 3-alpha-hydroxysteroid dehydrogenase activity. Has broad substrate specificity and acts on various 17-alpha-hydroxysteroids, 17-ketosteroids, 3-alpha hydroxysteroids and 3-ketosteroids. Reduction of keto groups is strictly stereoselective. Reduction of 17-ketosteroids yields only 17-alpha-hydroxysteroids. Likewise, reduction of 3-ketosteroids yields only 3-alpha-hydroxysteroids (By similarity).[UniProtKB/Swiss-Prot Function]