

# Product datasheet for RR202092L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Wee1 (NM\_001012742) Rat Tagged ORF Clone Lentiviral Particle

### **Product data:**

Product Type: Lentiviral Particles

**Product Name:** Wee1 (NM\_001012742) Rat Tagged ORF Clone Lentiviral Particle

Symbol: Wee1

Synonyms: MGC105683

Mammalian Cell

Puromycin

Selection:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001012742

ORF Size: 1938 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RR202092).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001012742.1

 RefSeq Size:
 3305 bp

 RefSeq ORF:
 1941 bp

 Locus ID:
 308937

 UniProt ID:
 Q63802

Cytogenetics: 1q33







### **Gene Summary:**

Acts as a negative regulator of entry into mitosis (G2 to M transition) by protecting the nucleus from cytoplasmically activated cyclin B1-complexed CDK1 before the onset of mitosis by mediating phosphorylation of CDK1 on 'Tyr-15'. Specifically phosphorylates and inactivates cyclin B1-complexed CDK1 reaching a maximum during G2 phase and a minimum as cells enter M phase. Phosphorylation of cyclin B1-CDK1 occurs exclusively on 'Tyr-15' and phosphorylation of monomeric CDK1 does not occur. Its activity increases during S and G2 phases and decreases at M phase when it is hyperphosphorylated. A correlated decrease in protein level occurs at M/G1 phase, probably due to its degradation (By similarity). [UniProtKB/Swiss-Prot Function]