

# Product datasheet for RR202254L4V

# Csk (NM\_001030039) Rat Tagged ORF Clone Lentiviral Particle

## **Product data:**

#### **Product Type: Lentiviral Particles Product Name:** Csk (NM\_001030039) Rat Tagged ORF Clone Lentiviral Particle Symbol: Csk MGC112926 Synonyms: **Mammalian Cell** Puromycin Selection: Vector: pLenti-C-mGFP-P2A-Puro (PS100093) mGFP Tag: NM 001030039 ACCN: ORF Size: 1350 bp The ORF insert of this clone is exactly the same as(RR202254). **ORF** Nucleotide Sequence: The molecular sequence of this clone aligns with the gene accession number as a point of **OTI Disclaimer:** reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info **OTI** Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. **RefSeq:** NM 001030039.1, NP 001025210.1 **RefSeq Size:** 2108 bp **RefSeq ORF:** 1353 bp Locus ID: 315707 **UniProt ID:** P32577 Cytogenetics: 8q24



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### 

Gene Summary:Non-receptor tyrosine-protein kinase that plays an important role in the regulation of cell<br/>growth, differentiation, migration and immune response. Phosphorylates tyrosine residues<br/>located in the C-terminal tails of Src-family kinases (SFKs) including LCK, SRC, HCK, FYN, LYN,<br/>CSK or YES1. Upon tail phosphorylation, Src-family members engage in intramolecular<br/>interactions between the phosphotyrosine tail and the SH2 domain that result in an inactive<br/>conformation. To inhibit SFKs, CSK is recruited to the plasma membrane via binding to<br/>transmembrane proteins or adapter proteins located near the plasma membrane.<br/>Suppresses signaling by various surface receptors, including T-cell receptor (TCR) and B-cell<br/>receptor (BCR) by phosphorylating and maintaining inactive several positive effectors such as<br/>FYN or LCK (By similarity).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US